

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
4 October 2001 (04.10.2001)

PCT

(10) International Publication Number  
WO 01/72977 A2

- (51) International Patent Classification: C12N 9/00, 15/52, C07K 16/40, A01K 67/027, C12N 15/66, C12Q 1/68, G01N 33/53, A61K 38/43
- (21) International Application Number: PCT/US01/09659
- (22) International Filing Date: 27 March 2001 (27.03.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/192,408 27 March 2000 (27.03.2000) US  
60/212,725 20 June 2000 (20.06.2000) US  
09/609,816 3 July 2000 (03.07.2000) US
- (71) Applicant: PE CORPORATION (NY) [US/US]; Robert A. Millman, Assistant Secretary, 761 Main Avenue, Norwalk, CT 06859 (US).
- (72) Inventors: WOODAGE, Trevor; Celera, 45 West Gude Drive, Rockville, MD 20850 (US). WEI, Ming-Hui; Celera, 45 West Gude Drive, Rockville, MD 20850 (US). KODIRA, Chinnappa; Celera, 45 West Gude Drive, Rockville, MD 20850 (US). DI FRANCESCO, Valentina; Celera, 45 West Gude Drive, Rockville, MD 20850 (US). BEASLEY, Ellen, M.; Celera, 45 West Gude Drive, Rockville, MD 20850 (US).
- (74) Agent: MILLMAN, Robert, A.; Celera Genomics Corp., 45 West Gude Drive C2-4, Rockville, MD 20850 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

[Continued on next page]

(54) Title: ISOLATED HUMAN DRUG-METABOLIZING PROTEINS, NUCLEIC ACID MOLECULES ENCODING HUMAN DRUG-METABOLIZING PROTEINS, AND USES THEREOF

(57) Abstract: The present invention provides amino acid sequences of peptides that are encoded by genes within the human genome, the proteins of the present invention. The present invention specifically provides isolated peptide and nucleic acid molecules, methods of identifying orthologs and paralogs of the proteins of the present invention, and methods of identifying modulators of the proteins of the present invention.

<110> Woodage, Trevor  
Wei, Ming-Hui  
Kodira, Chinnappa  
Beasley, Ellen  
DiFrancesco, Valentina

<120> ISOLATED HUMAN DRUG-METABOLIZING  
PROTEINS, NUCLEIC ACID MOLECULES ENCODING HUMAN  
DRUG-METABOLIZING PROTEINS, AND USES THEREOF

<130> CL000669PCT

<140> N/A  
<141> 2001-0327

<150> 60/192,408  
<151> 2000-03-27

<150> 60/212,725  
<151> 2000-06-20

<150> 09/609,816  
<151> 2000-07-03

<160> 23

<170> FASTSEQ for Windows Version 4.0

<210> 1  
<211> 942  
<212> DNA  
<213> HUMAN

<400> 1  
atctccatgt gctgagatgt agaaacacgc tccacagatg gaaacaaagc cagacttgtt 60  
taacatcatg gactgagatg gactccatgc gttgatatta tcaaaagatc ggtggaagaa 120  
agatgtgtaat ttcaaaagca agctgagatg tcttattctg gaaattatcc caaagtccag 180  
tacaacatgt atgcatgaaa tttaagatcat gattctaatat gatgtgagtg tggagaaatt 240  
caaaagagac cagactctag atagacacgc ttctcttgaa ctgaatttcc cccataaaga 300  
aaacatgatc ttgagatctg tcttgaaatc gttctcagca caactgatac aaacatctct 360  
ccttcacatc atgattccac catctatctg gaaagaaac tgaagatctg tctatgtgga 420  
cagaatccc aagatttgcg tgggtctata ctaccacttt cagagatgag cttcttctat 480  
gctgtatctc cagactttag agaaatttta tgaagatttc atgtccgaga aagttgttgg 540  
cgggtccatg ttgagatctg tgaagagatg gttgttgcga aaagacatgc accgatctct 600  
ctactctctc taaagagata ttaaaasaaa taaaacatc gaagatccaa aggtgttga 660  
atctctgaga aaacttgggt caggtgatgt taaaacaaag atgtccacac atactctatc 720  
tgatgtatgt aagataatc ccatggccaa ccatctatgc gtaactgttc acatattcaa 780  
tcaactatcc tcaaaatta tgaagaaatg gactctgaga gactgtgaga accatttcaa 840  
tgtgtttgtg aatgaaact ttgataagca tatgaaag aagatggcag ggtccacact 900  
gaactctgct ctgagatctc gagagaaaca caacaaactc ag 942

<210> 2  
<211> 922  
<212> DNA  
<213> HUMAN

<400> 2  
atctccatgt gctgagatgt agaaacacgc tccacagatg gaaacaaagc cagacttgtt 60  
taacatcatg gactgagatg gactccatgc gttgatatta tcaaaagatc ggtggaagaa 120  
agatgtgtaat ttcaaaagca agctgagatg tcttattctg gaaattatcc caaagtccag 180  
tacaacatgt atgcatgaaa tttaagatcat gattctaatat gatgtgagtg tggagaaatt 240

WO 01/72977 A2

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**ISOLATED HUMAN DRUG-METABOLIZING PROTEINS, NUCLEIC ACID  
MOLECULES ENCODING HUMAN DRUG-METABOLIZING PROTEINS,  
AND USES THEREOF**

5       **RELATED APPLICATIONS**

The present application claims priority to U.S. Serial Nos. 60/192,408, filed March 27, 2000 (Atty. Docket CL000399-PROV), 60/212,725, filed June 20, 2000 (Atty. Docket CL000669-PROV), and 09/609,816, filed July 3, 2000 (Atty. Docket CL000669).

10       **FIELD OF THE INVENTION**

The present invention is in the field of proteins that are related to the sulfotransferase drug-metabolizing enzyme subfamily, recombinant DNA molecules and protein production. The present invention specifically provides novel drug-metabolizing peptides and proteins and nucleic acid molecules encoding such protein molecules, for use in labeling reagents with tagged sulfur and for  
15 the development of human therapeutics and human therapeutic development.

**BACKGROUND OF THE INVENTION**

Drug-Metabolizing Proteins

20       Induction of drug-metabolizing enzymes ("DMEs") is a common biological response to xenobiotics, the mechanisms and consequences of which are important in academic, industrial, and regulatory areas of pharmacology and toxicology.

For most drugs, drug-metabolizing enzymes determine how long and how much of a drug remains in the body. Thus, developers of drugs recognize the importance of characterizing a drug  
25 candidate's interaction with these enzymes. For example, polymorphisms of the drug-metabolizing enzyme CYP2D6, a member of the cytochrome p450 ("CYP") superfamily, yield phenotypes of slow or ultra-rapid metabolizers of a wide spectrum of drugs including antidepressants, antipsychotics, beta-blockers, and antiarrhythmics. Such abnormal rates of drug metabolism can lead to drug ineffectiveness or to systemic accumulation and toxicity.

30       For pharmaceutical scientists developing a candidate drug, it is important know as early as possible in the design phase which enzymes metabolize the drug candidate and the speed with which they do it. Historically, the enzymes on a drug's metabolic pathway were determined through metabolism studies in animals, but this approach has now been largely supplanted by the use of human tissues or cloned drug-metabolizing enzymes to provide insights into the specific role of  
35 individual forms of these enzymes. Using these tools, the qualitative and quantitative fate of a drug

candidate can be predicted prior to its first administration to humans. As a consequence, the selection and optimization of desirable characteristics of metabolism are possible early in the development process, thus avoiding unanticipated toxicity problems and associated costs subsequent to the drug's clinical investigation. Moreover, the effect of one drug on another's disposition can be inferred.

Known drug-metabolizing enzymes include the cytochrome p450 ("CYP") superfamily, N-acetyl transferases ("NAT"), UDP-glucuronosyl transferases ("UGT"), methyl transferases, alcohol dehydrogenase ("ADH"), aldehyde dehydrogenase ("ALDH"), dihydropyrimidine dehydrogenase ("DPD"), NADPH:quinone oxidoreductase ("NQO" or "DT diaphorase"), catechol O-methyltransferase ("COMT"), glutathione S-transferase ("GST"), histamine methyltransferase ("HMT"), sulfotransferases ("ST"), thiopurine methyltransferase ("TPMT"), and epoxide hydroxylase. Drug-metabolizing enzymes are generally classified into two phases according to their metabolic function. Phase I enzymes catalyze modification of functional groups, and phase II enzymes catalyze conjugation with endogenous substituents. These classifications should not be construed as exclusive nor exhaustive, as other mechanisms of drug metabolism have been discovered. For example, the use of active transport mechanisms been characterized as part of the process of detoxification.

Phase I reactions include catabolic processes such as deamination of aminases, hydrolysis of esters and amides, conjugation reactions with, for example, glycine or sulfate, oxidation by the cytochrome p450 oxidation/reduction enzyme system and degradation in the fatty acid pathway. Hydrolysis reactions occur mainly in the liver and plasma by a variety of non-specific hydrolases and esterases. Both deaminases and amidases, also localized in the liver and serum, carry out a large part of the catabolic process. Reduction reactions occur mainly intracellularly in the endoplasmic reticulum.

Phase II enzymes detoxify toxic substances by catalyzing their conjugation with water-soluble substances, thus increasing toxins' solubility in water and increasing their rate of excretion. Additionally, conjugation reduces the toxins' biological reactivity. Examples of phase II enzymes include glutathione S-transferases and UDP-glucuronosyl transferases, which catalyze conjugation to glutathione and glucuronic acid, respectively. Transferases perform conjugation reactions mainly in the kidneys and liver.

The liver is the primary site of elimination of most drugs, including psychoactive drugs, and contains a plurality of both phase I and phase II enzymes that oxidize or conjugate drugs, respectively.



Physicians currently prescribe drugs and their dosages based on a population average and fail to take genetic variability into account. The variability between individuals in drug metabolism is usually due to both genetic and environmental factors, in particular, how the drug-metabolizing enzymes are controlled. With certain enzymes, the genetic component predominates and variability is associated with variants of the normal, wild-type enzyme.

Most drug-metabolizing enzymes exhibit clinically relevant genetic polymorphisms. Essentially all of the major human enzymes responsible for modification of functional groups or conjugation with endogenous substrates exhibit common polymorphisms at the genomic level. For example, polymorphisms expressing a non-functioning variant enzyme results in a sub-group of patients in the population who are more prone to the concentration-dependent effects of a drug. This sub-group of patients may show toxic side effects to a dose of drug that is otherwise without side effects in the general population. Recent development in genotyping allows identification of affected individuals. As a result, their atypical metabolism and likely response to a drug metabolized by the affected enzyme can be understood and predicted, thus permitting the physician to adjust the dose of drug they receive to achieve improved therapy.

A similar approach is also becoming important in identifying risk factors associated with the development of various cancers. This is because the enzymes involved in drug metabolism are also responsible for the activation and detoxification of chemical carcinogens. Specifically, the development of neoplasia is regulated by a balance between phase I enzymes, which activate carcinogens, and phase II enzymes, which detoxify them. Accordingly, an individual's susceptibility to cancer often involves the balance between these two processes, which is, in part, genetically determined and can be screened by suitable genotyping tests. Higher induction of phase I enzymes compared to phase II enzymes results in the generation of large amounts of electrophiles and reactive oxygen species and may cause DNA and membrane damage and other adverse effects leading to neoplasia. Conversely, higher levels of phase II enzyme expression can protect cells from various chemical compounds.

Abnormal activity of drug-metabolizing enzymes has been implicated in a range of human diseases, including cancer, Parkinson's disease, myotonic dystrophy, and developmental defects.

### Cytochrome p450

An example of a phase I drug-metabolizing enzyme is the cytochrome p450 ("CYP") superfamily, the members of which comprise the major drug-metabolizing enzymes expressed in the liver. The CYP superfamily comprises heme proteins which catalyze the oxidation and dehydrogenation of a number of endogenous and exogenous lipophilic compounds. The CYP

superfamily has immense diversity in its functions, with hundreds of isoforms in many species catalyzing many types of chemical reactions. The CYP superfamily comprises at least 30 related enzymes, which are divided into different families according to their amino acid homology. Examples of CYP families include CYP families 1, 2, 3 and 4, which comprise endoplasmic

5 reticulum proteins responsible for the metabolism of drugs and other xenobiotics. Approximately 10-15 individual gene products within these four families metabolize thousands of structurally diverse compounds. It is estimated that collectively the enzymes in the CYP superfamily participate in the metabolism of greater than 80% of all available drugs used in humans. For example, the CYP 1A subfamily comprises CYP 1A2, which metabolizes several widely used drugs, including

10 acetaminophen, amitriptyline, caffeine, clozapine, haloperidol, imipramine, olanzapine, ondansetron, phenacetin, propafenone, propranolol, tacrine, theophylline, verapamil. In addition, CYP enzymes play additional roles in the metabolism of some endogenous substrates including prostaglandins and steroids.

Some CYP enzymes exist in a polymorphic form, meaning that a small percentage of the

15 population possesses mutant genes that alter the activity of the enzyme, usually by diminishing or abolishing activity. For example, a genetic polymorphism has been well characterized with the CYP 2C19 and CYP 2D6 genes. Substrates of CYP 2C19 include clomipramine, diazepam, imipramine, mephenytoin, moclobemide, omeprazole, phenytoin, propranolol, and tolbutamide. Substrates of CYP 2D6 include alprenolol, amitriptyline, chlorpheniramine, clomipramine, codeine, desipramine,

20 dextromethorphan, encainide, fluoxetine, haloperidol, imipramine, indoramin, metoprolol, nortriptyline, ondansetron, oxycodone, paroxetine, propranolol, and propafenone. Polymorphic variants of these genes metabolize these substrates at different rates, which can effect a patient's effective therapeutic dosage.

While the substrate specificity of CYPs must be very broad to accommodate the metabolism

25 of all of these compounds, each individual CYP gene product has a narrower substrate specificity defined by its binding and catalytic sites. Drug metabolism can thereby be regulated by changes in the amount or activity of specific CYP gene products. Methods of CYP regulation include genetic differences in the expression of CYP gene products (i.e., genetic polymorphisms), inhibition of CYP metabolism by other xenobiotics that also bind to the CYP, and induction of certain CYPs by

30 the drug itself or other xenobiotics. Inhibition and induction of CYPs is one of the most common mechanisms of adverse drug interactions. For example, the CYP3A subfamily is involved in clinically significant drug interactions involving nonsedating antihistamines and cisapride that may result in cardiac dysrhythmias. In another example, CYP3A4 and CYP1A2 enzymes are involved in drug interactions involving theophylline. In yet another example, CYP2D6 is responsible for the

metabolism of many psychotherapeutic agents. Additionally, CYP enzymes metabolize the protease inhibitors used to treat patients infected with the human immunodeficiency virus. By understanding the unique functions and characteristics of these enzymes, physicians may better anticipate and manage drug interactions and may predict or explain an individual's response to a particular therapeutic regimen.

Examples of reactions catalyzed by the CYP superfamily include peroxidative reactions utilizing peroxides as oxygen donors in hydroxylation reactions, as substrates for reductive beta-scission, and as peroxyhemiacetal intermediates in the cleavage of aldehydes to formate and alkenes. Lipid hydroperoxides undergo reductive beta-cleavage to give hydrocarbons and aldehydic acids. One of these products, trans-4-hydroxynonenal, inactivates CYP, particularly alcohol-inducible 2E1, in what may be a negative regulatory process. Although a CYP iron-oxene species is believed to be the oxygen donor in most hydroxylation reactions, an iron-peroxy species is apparently involved in the deformylation of many aldehydes with desaturation of the remaining structure, as in aromatization reactions.

Examples of drugs with oxidative metabolism associated with CYP enzymes include acetaminophen, alfentanil, alprazolam, alprenolol, amiodarone, amitriptyline, astemizole, buspirone, caffeine, carbamazepine, chlorpheniramine, cisapride, clomipramine, clomipramine, clozapine, codeine, colchicine, cortisol, cyclophosphamide, cyclosporine, dapsone, desipramine, dextromethorphan, diazepam, diclofenac, diltiazem, encainide, erythromycin, estradiol, felodipine, fluoxetine, fluvastatin, haloperidol, ibuprofen, imipramine, indinavir, indomethacin, indoramin, irbesartan, lidocaine, losartan, macrolide antibiotics, mephenytoin, methadone, metoprolol, mexilitene, midazolam, moclobemide, naproxen, nefazodone, nifedipine, nifedipine, nitrendipine, nortriptyline, olanzapine, omeprazole, ondansetron, oxycodone, paclitaxel, paroxetine, phenacetin, phenytoin, piroxicam, progesterone, propafenone, propranolol, quinidine, ritonavir, saquinavir, sertraline, sildenafil, S-warfarin, tacrine, tamoxifen, tenoxicam, terfenadine, testosterone, theophylline, timolol, tolbutamide, triazolam, verapamil, and vinblastine.

Abnormal activity of phase I enzymes has been implicated in a range of human diseases. For example, enhanced CYP2D6 activity has been related to malignancies of the bladder, liver, pharynx, stomach and lungs, whereas decreased CYP2D activity has been linked to an increased risk of Parkinson's disease. Other syndromes and developmental defects associated with deficiencies in the CYP superfamily include cerebrotendinous xanthomatosis, adrenal hyperplasia, gynecomastia, and myotonic dystrophy.

The CYP superfamily are a major target for drug action and development. Accordingly, it is valuable to the field of pharmaceutical development to identify and characterize previously unknown members of the CYP superfamily.

5        UDP-glucuronosyltransferases

Potential drug interactions involving phase II metabolism are increasingly being recognized. An important group of phase II enzymes involved in drug metabolism are the glucuronosyltransferases, especially the UDP-glucuronyltransferase ("UGT") superfamily. Members of the UGT superfamily catalyze the enzymatic addition of UDP glucuronic acid as a sugar donor to fat-soluble chemicals, a process which increases their solubility in water and increases their rate of excretion. In mammals, glucuronic acid is the main sugar that is used to prevent the accumulation of waste products of metabolism and fat-soluble chemicals from the environment to toxic levels in the body. Both inducers and inhibitors of glucuronosyltransferases are known and have the potential to affect the plasma concentration and actions of important drugs, including psychotropic drugs.

The UGT superfamily comprises several families of enzymes in several species defined with a nomenclature similar to that used to define members of the CYP superfamily. In animals, yeast, plants and bacteria there are at least 110 distinct known members of the UGT superfamily. As many as 33 families have been defined, with three families identified in humans. Different UGT families are defined as having <45% amino acid sequence homology; within subfamilies there is approximately 60% homology. The members of the UGT superfamily are part of a further superfamily of UDP glycosyltransferases found in animals, plants and bacteria.

The role of phase II enzymes, and of UGT enzymes in particular, is being increasingly recognized as important in psychopharmacology. UGT enzymes conjugate many important psychotropic drugs and are an important source of variability in drug response and drug interactions. For example, the benzodiazepines lorazepam, oxazepam, and temazepam undergo phase II reactions exclusively before being excreted into the urine.

Phase II enzymes metabolize and detoxify hazardous substances, such as carcinogens. The expression of genes encoding phase II enzymes is known to be up-regulated by hundreds of agents. For example, oltipraz is known to up-regulate phase II enzyme expression. Studies have demonstrated protection from the cancer-causing effects of carcinogens when selected phase II enzyme inducers are administered prior to the carcinogens. The potential use of phase II enzyme inducers in humans for prevention of cancers related to exposure to carcinogens has prompted studies aimed at understanding their molecular effects. Current biochemical and molecular

biological research methodologies can be used to identify and characterize selective phase II enzyme inducers and their targets. Identification of genes responding to cancer chemopreventive agents will facilitate studies of their basic mechanism and provide insights about the relationship between gene regulation, enzyme polymorphism, and carcinogen detoxification.

5        Examples of drugs with conjugative metabolism associated with UGT enzymes include amitriptyline, buprenorphine, chlorpromazine, clozapine, codeine, cyproheptadine, dihydrocodeine, doxepin, imipramine, lamotrigine, lorazepam, morphine, nalorphine, naltrexone, temazepam, and valproate.

Abnormal activity of phase II enzymes has been implicated in a range of human diseases.  
10    For example, Gilbert syndrome is an autosomal dominant disorder caused by mutation in the UGT1 gene, and mutations in the UGT1A1 enzyme have been demonstrated to be responsible for Crigler-Najjar syndrome.

The UGT superfamily are a major target for drug action and development. Accordingly, it is valuable to the field of pharmaceutical development to identify and characterize previously unknown  
15    members of the UGT superfamily.

#### Sulfotransferases

Sulfation and sulfate conjugate hydrolysis play an important role in metabolism, and are catalysed by members of the sulfotransferase and sulfatase enzyme super-families.

20    Sulfotransferases are enzymes that transfer sulfate groups to acceptor molecules. These enzymes are involved in the posttranslational sulfation of proteins and the sulfate conjugation of exogenous chemicals and bile acids. Cytosolic sulfotransferases in mammalian cells catalyze the transfer of a sulfonate group from 3'-phosphoadenosine- 5'-phosphosulfate, the active sulfate, to the hydroxyl groups or amine groups of substrate compounds. Traditionally, they are viewed as detoxifying or  
25    Phase II drug-metabolizing enzymes that facilitate the removal of drugs and xenobiotic compounds. Increasingly, however, these enzymes have been shown to be involved in the sulfation of endogenous compounds, such as thyroid and steroid hormones, catecholamines, and bile acids, to fulfill fundamental biochemical/physiological needs. The balance of sulfoconjugation (by sulfotransferases) and deconjugation (by sulfotases) may have physiological implications; in  
30    addition to catecholamine release, it may determine the availability of free catecholamines during diurnal rhythms and stress or modify their renal excretion. Circumstantial evidence, including a close homology within the aryl sulfatases and steroid sulfatase gene, the first implicated in catecholamine metabolism, the second in steroid metabolism, suggests a genetic defect of sulfatases

in essential hypertension. A similar, but secondary, sulfatase defect may affect catecholamine metabolism and action in chronic renal failure.

In general, sulfation is a deactivating, detoxication pathway, but for some chemicals the sulfate conjugates are much more reactive than the parent compound (Coughtrie MW, et al., 1998.

5 Biology and function of the reversible sulfation pathway catalysed by human sulfotransferases and sulfatases. *Chem Biol Interact.* 20:109(1-3):3-27). For example N-hydroxyarylamines sulfotransferase (HAST-I) detoxifies phenols but activates N-hydroxyarylamines into a mutagenic form, N-hydroxy-2-acetylaminofluorene (Nagata, K., et al., 1993. Isolation and expression of a cDNA encoding a male-specific rat sulfotransferase that catalyzes activation of N-hydroxy-2-

10 acetylaminofluorene. *J. Biol. Chem.* 268 (33), 24720-24725 and Gong DW, et al., 1991. Purification of hepatic N-hydroxyarylamines sulfotransferases and their regulation by growth hormone and thyroid hormone in rats. *J Biochem (Tokyo)* 1991 Aug;110(2):226-31.).

The discovery of proteins that are related to and are potentially new sulfotransferase satisfies a need in the art by providing new compositions which are useful towards the design of stable and

15 more active drugs, and in the prevention, diagnosis, and treatment of cancer, hypertension, renal diseases, and neurodegenerative diseases.

In addition to such use in the study of drug metabolism, sulfotransferases are also useful in labeling agents with tagged sulfur. This provides another use of the proteins of the present invention.

20 Further, since the genes of the present invention are mapped to chromosomal position and SNP variation is present, the nucleic acid molecules of the present invention can be used in linkage studies and genetic mapping uses.

#### SUMMARY OF THE INVENTION

25 The present invention is based in part on the identification of amino acid sequences of proteins that are related to the sulfotransferase drug-metabolizing enzyme subfamily, as well as allelic variants and other mammalian orthologs thereof. These unique protein sequences, and nucleic acid sequences that encode these proteins, can be used to label proteins, as models for the development of human therapeutic targets, aid in the identification of therapeutic proteins, and

30 serve as targets for the development of human therapeutic agents that modulate protein of the present invention activity in cells and tissues that express the protein of the present invention. Specially, experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine. Further, the nucleic acid molecules herein described can be used to label agents with tagged sulfur.

## DESCRIPTION OF THE FIGURE SHEETS

FIGURE 1 provides the nucleotide sequence of two alternative transcript sequences (SEQ ID NOS:1 and 2) and a cDNA sequence (SEQ ID NO:3) that encodes the protein of the present invention. In addition, structure and functional information is provided, such as ATG start, stop and tissue distribution, that allows one to readily determine specific uses of inventions based on this molecular sequence. Experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine.

FIGURE 2 provides the amino acid sequences of the proteins of the present invention. (SEQ ID NOS:4-6) In addition, structural and functional information such as protein family, function, important domains and modification sites is provided, allowing one to readily determine specific uses of inventions based on this molecular sequence.

FIGURE 3 provides genomic sequences that span the gene encoding the protein of the present invention. (SEQ ID NO:7) In addition structural and functional information, such as intron/exon structure, promoter location, etc., is provided allowing one to readily determine specific uses of inventions based on this molecular sequence. For example, as illustrated in Figure 3, identified SNP variations include a1008c, a6094g, t7898c, a10018c, a10791t, a12451t, t14674c, g14860c, a20133g, a1025c, t12602c, g12829t, a1442t.

## DETAILED DESCRIPTION OF THE INVENTION

### General Description

The present invention is based on the gene identification that occurred during the sequencing of the human genome. During the sequencing and assembly of the human genome, analysis of the sequence information revealed previously unidentified fragments of the human genome that encode peptides that share structural and/or sequence homology to protein/peptide/domains identified and characterized within the art as being a drug-metabolizing enzyme protein or part of a drug-metabolizing enzyme protein and are specifically related to the sulfotransferase drug-metabolizing enzyme subfamily. Utilizing these sequences, additional genomic sequences were assembled and transcript and/or cDNA sequences were isolated and characterized. Based on this analysis, the present invention provides amino acid sequences of proteins that are related to the sulfotransferase drug-metabolizing enzyme subfamily, nucleic acid sequences in the form of transcript sequences, cDNA sequences and/or genomic sequences that encode these proteins, nucleic acid variation (allelic information), tissue distribution of expression.

and information about the closest art known protein/peptide/domain that has structural or sequence homology to the protein of the present invention.

In addition to being previously unknown, the proteins that are provided in the present invention are selected based on their ability to be used for the development of commercially important products and services. Specifically, the present proteins are selected based on homology and/or structural relatedness to known drug-metabolizing enzyme proteins of the sulfotransferase drug-metabolizing enzyme subfamily and the expression pattern observed. Specifically, experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine. The art has clearly established the commercial importance of members of this family of proteins and proteins that have expression patterns similar to that of the present gene for industrial uses, such as labeling agents with tagged sulfur, genetic mapping uses, use of the SNP and chromosomal position for linkage studies, genetic expression uses, use of the control region to drive expression in the kidneys, etc. Some of the more specific features of the proteins of the present invention, and the uses thereof, are described herein, particularly in the Background of the Invention and in the annotation provided in the Figures, and/or are known within the art for the known sulfotransferase subfamily of drug-metabolizing enzyme proteins.

#### Specific Embodiments

##### Peptide Molecules

The present invention provides nucleic acid sequences that encode protein molecules that have been identified as being related to the sulfotransferase drug-metabolizing enzyme subfamily (protein sequences are provided in Figure 2, transcript/cDNA sequences are provided in Figure 1 and genomic sequences are provided in Figure 3). The peptide sequences provided in Figure 2, as well as the obvious variants described herein, particularly allelic variants as identified herein and using the information in Figure 3, will be referred herein as the proteins of the present invention, or proteins of the present invention.

The present invention provides isolated peptide and protein molecules that consist of, consist essentially of, or comprise the amino acid sequences of the proteins disclosed in the Figure 2, (encoded by the nucleic acid molecule shown in Figure 1, transcript/cDNA or Figure 3, genomic sequence), as well as all obvious variants of these peptides that are within the art to make and use. Some of these variants are described in detail below.

As used herein, a peptide is said to be "isolated" or "purified" when it is substantially free of cellular material or free of chemical precursors or other chemicals. The proteins of the present



invention can be purified to homogeneity or other degrees of purity. The level of purification will be based on the intended use. The critical feature is that the preparation allows for the desired function of the peptide, even if in the presence of considerable amounts of other components (the features of an isolated nucleic acid molecule is discussed below).

5 In some uses, "substantially free of cellular material" includes preparations of the peptide having less than about 30% (by dry weight) other proteins (i.e., contaminating protein), less than about 20% other proteins, less than about 10% other proteins, or less than about 5% other proteins. When the peptide is recombinantly produced, it can also be substantially free of culture medium, i.e., culture medium represents less than about 20% of the volume of the protein preparation.

10 The language "substantially free of chemical precursors or other chemicals" includes preparations of the peptide in which it is separated from chemical precursors or other chemicals that are involved in its synthesis. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of the peptide having less than about 30% (by dry weight) chemical precursors or other chemicals, less than about 20% chemical precursors or other  
15 chemicals, less than about 10% chemical precursors or other chemicals, or less than about 5% chemical precursors or other chemicals.

The isolated protein can be purified from cells that naturally express it, purified from cells that have been altered to express it (recombinant), or synthesized using known protein synthesis methods. Experimental data provided in Figure\*1 shows that the sulfotransferase of the present invention is  
20 expressed at least in the kidney and small intestine. For example, a nucleic acid molecule encoding the proteins of the present invention is cloned into an expression vector, the expression vector introduced into a host cell and the protein expressed in the host cell. The protein can then be isolated from the cells by an appropriate purification scheme using standard protein purification techniques. Many of these techniques are described in detail below.

25 Accordingly, the present invention provides proteins that consist of the amino acid sequences provided in Figure 2 (SEQ ID NOS:4-6), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NOS:1 and 2, alternative transcript sequences, and SEQ ID NO:3, cDNA sequences) and the genomic sequences provided in Figure 3 (SEQ ID NO:7). The amino acid sequence of such a protein is provided in Figure 2. A protein consists of an amino acid  
30 sequence when the amino acid sequence is the final amino acid sequence of the protein. Such a protein can further comprise the variants provided in Figure 3.

The present invention further provides proteins that consist essentially of the amino acid sequences provided in Figure 2 (SEQ ID NOS:4-6), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NOS:1 and 2, alternative

transcript sequences, and SEQ ID NO:3, cDNA sequences) and the genomic sequences provided in Figure 3 (SEQ ID NO:7). A protein consists essentially of an amino acid sequence when such an amino acid sequence is present with only a few additional amino acid residues, for example from about 1 to about 100 or so additional residues, typically from 1 to about 20 additional residues in the final protein.

The present invention further provides proteins that comprise the amino acid sequences provided in Figure 2 (SEQ ID NOS:4-6), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NOS:1 and 2, alternative transcript sequences, and SEQ ID NO:3, cDNA sequences) and the genomic sequences provided in Figure 3 (SEQ ID NO:7). A protein comprises an amino acid sequence when the amino acid sequence is at least part of the final amino acid sequence of the protein. In such a fashion, the protein can be only the peptide or have additional amino acid molecules, such as amino acid residues (contiguous encoded sequence) that are naturally associated with it or heterologous amino acid residues/peptide sequences. Such a protein can have a few additional amino acid residues or can comprise several hundred or more additional amino acids. The preferred classes of proteins that are comprised of the proteins of the present invention are the naturally occurring mature proteins. A brief description of how various types of these proteins can be made/isolated is provided below.

The proteins of the present invention can be attached to heterologous sequences to form chimeric or fusion proteins. Such chimeric and fusion proteins comprise one of the proteins of the present invention operatively linked to a heterologous protein having an amino acid sequence not substantially homologous to the proteins of the present invention. "Operatively linked" indicates that the proteins of the present invention and the heterologous protein are fused in-frame. The heterologous protein can be fused to the N-terminus or C-terminus of the proteins of the present invention.

In some uses, the fusion protein does not affect the activity of the proteins of the present invention *per se*. For example, the fusion protein can include, but is not limited to, enzymatic fusion proteins, for example beta-galactosidase fusions, yeast two-hybrid GAL fusions, poly-His fusions, MYC-tagged, HI-tagged and Ig fusions. Such fusion proteins, particularly poly-His fusions, can facilitate the purification of recombinant proteins of the present invention. In certain host cells (e.g., mammalian host cells), expression and/or secretion of a protein can be increased by using a heterologous signal sequence.

A chimeric or fusion protein can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different protein sequences are ligated together in-frame in accordance with conventional techniques. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification

of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and re-amplified to generate a chimeric gene sequence (see Ausubel *et al.*, *Current Protocols in Molecular Biology*, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST protein). A protein of the present invention-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the proteins of the present invention.

As mentioned above, the present invention also provides and enables obvious variants of the amino acid sequence of the proteins of the present invention, such as naturally occurring mature forms of the peptide, allelic/sequence variants of the peptides, non-naturally occurring recombinantly derived variants of the peptides, and orthologs and paralogs of the peptides. Such variants can readily be generated using art-known techniques in the fields of recombinant nucleic acid technology and protein biochemistry. It is understood, however, that variants exclude any amino acid sequences disclosed prior to the invention.

Such variants can readily be identified/made using molecular techniques and the sequence information disclosed herein. Further, such variants can readily be distinguished from other peptides based on sequence and/or structural homology to the proteins of the present invention. The degree of homology/identity present will be based primarily on whether the peptide is a functional variant or non-functional variant, the amount of divergence present in the paralog family and the evolutionary distance between the orthologs.

To determine the percent identity of two amino acid sequences or two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, at least 30%, 40%, 50%, 60%, 70%, 80%, or 90% or more of the length of a reference sequence is aligned for comparison purposes. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity and similarity between two sequences can be accomplished using a mathematical algorithm. (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part 1*, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch (*J. Mol. Biol.* (48):444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossom 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (Devereux, J., *et al.*, *Nucleic Acids Res.* 12(1):387 (1984)) (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. In another embodiment, the percent identity between two amino acid or nucleotide sequences is determined using the algorithm of E. Myers and W. Miller (CABIOS, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

The nucleic acid and protein sequences of the present invention can further be used as a "query sequence" to perform a search against sequence databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* (*J. Mol. Biol.* 215:403-10 (1990)). BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to the nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the proteins of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.* (*Nucleic Acids Res.* 25(17):3389-3402 (1997)). When utilizing BLAST and gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used.

Full-length pre-processed forms, as well as mature processed forms, and splice variants proteins that comprise one of the proteins of the present invention can readily be identified as having complete sequence identity to at least a portion of one of the proteins of the present invention as well as

being encoded by the same genetic locus as the proteins of the present invention provided herein. Figure 3, the map position was determined to be on chromosome 2 near marker SHGC-1639 (LOD score 10.01) using radiation hybrid mapping.

Allelic variants of a protein of the present invention can readily be identified as being a human  
5 protein having a high degree (significant) of sequence homology/identity to at least a portion of the proteins of the present invention as well as being encoded by the same genetic locus as the proteins of the present invention provided herein. Genetic locus can readily be determined based on the genomic information provided in Figure 3, such as the genomic sequence mapped to the reference human. As indicated by the data presented in Figure 3, the map position was determined to be on chromosome 2  
10 near marker SHGC-1639 (LOD score 10.01) using radiation hybrid mapping. As used herein, two proteins (or a region of the proteins) have significant homology when the amino acid sequences are typically at least about 70-80%, 80-90%, and more typically at least about 90-95% or more homologous. A significantly homologous amino acid sequence, according to the present invention, will be encoded by a nucleic acid sequence that will hybridize to a protein of the present invention  
15 encoding nucleic acid molecule under stringent conditions as more fully described below.

Figure 3 provides SNP information that was identified from chromosomal sequence obtained from a number of individuals as well as SNPs found during the cDNA cloning process. These include, a6094g, t7898c, a10018c, a10791t, a12451t, t14674c, g14860c, a20133g, a1025c, t12602c, g12829t, a1442t.

20 Paralogs of a protein of the present invention can readily be identified as having some degree of significant sequence homology/identity to at least a portion of the proteins of the present invention, as being encoded by a gene from humans, and as having similar activity or function. Two proteins will typically be considered paralogs when the amino acid sequences are typically at least about 60% or greater, and more typically at least about 70% or greater homology through a given region  
25 or domain. Such paralogs will be encoded by a nucleic acid sequence that will hybridize to a protein of the present invention encoding nucleic acid molecule under moderate to stringent conditions as more fully described below.

Orthologs of a protein of the present invention can readily be identified as having some degree of significant sequence homology/identity to at least a portion of the proteins of the present invention  
30 as well as being encoded by a gene from another organism. Preferred orthologs will be isolated from mammals, preferably primates, for the development of human therapeutic targets and agents. Such orthologs will be encoded by a nucleic acid sequence that will hybridize to a protein of the present invention encoding nucleic acid molecule under moderate to stringent conditions, as more fully described below, depending on the degree of relatedness of the two organisms yielding the proteins.

Non-naturally occurring variants of the proteins of the present invention can readily be generated using recombinant techniques. Such variants include, but are not limited to deletions, additions and substitutions in the amino acid sequence of the proteins of the present invention. For example, one class of substitutions are conserved amino acid substitution. Such substitutions are those  
5 that substitute a given amino acid in a protein of the present invention by another amino acid of like characteristics. Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu, and Ile; interchange of the hydroxyl residues Ser and Thr; exchange of the acidic residues Asp and Glu; substitution between the amide residues Asn and Gln; exchange of the basic residues Lys and Arg; and replacements among the aromatic residues Phe  
10 and Tyr. Guidance concerning which amino acid changes are likely to be phenotypically silent are found in Bowie *et al.*, *Science* 247:1306-1310 (1990).

Variant proteins can be fully functional or can lack function in one or more activities, e.g. ability to bind substrate, ability to phosphorylate substrate, ability to mediate signaling, etc. Fully functional variants typically contain only conservative variation or variation in non-critical residues or  
15 in non-critical regions. Figure 2 provides the result of protein analysis and can be used to identify critical domains/regions. Functional variants can also contain substitution of similar amino acids that result in no change or an insignificant change in function. Alternatively, such substitutions may positively or negatively affect function to some degree.

Non-functional variants typically contain one or more non-conservative amino acid  
20 substitutions, deletions, insertions, inversions, or truncation or a substitution, insertion, inversion, or deletion in a critical residue or critical region.

Amino acids that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham *et al.*, *Science* 244:1081-1085 (1989)), particularly using the results provided in Figure 2. The latter procedure introduces single  
25 alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as drug-metabolizing enzyme activity or in assays such as an *in vitro* proliferative activity. Sites that are critical for binding partner/substrate binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith *et al.*, *J. Mol. Biol.* 224:899-904 (1992); de Vos *et al.* *Science* 255:306-312 (1992)).

30 The present invention further provides fragments of the proteins of the present invention, in addition to proteins and peptides that comprise and consist of such fragments, particularly those comprising the residues identified in Figure 2. The fragments to which the invention pertains, however, are not to be construed as encompassing fragments that may be disclosed publicly prior to the present invention.

As used herein, a fragment comprises at least 8, 10, 12, 14, 16, or more contiguous amino acid residues from a protein of the present invention. Such fragments can be chosen based on the ability to retain one or more of the biological activities of the proteins of the present invention or could be chosen for the ability to perform a function, e.g. bind a substrate or act as an immunogen. Particularly important fragments are biologically active fragments, peptides that are, for example, about 8 or more amino acids in length. Such fragments will typically comprise a domain or motif of the proteins of the present invention, e.g., active site, a transmembrane domain or a substrate-binding domain. Further, possible fragments include, but are not limited to, domain or motif containing fragments, soluble peptide fragments, and fragments containing immunogenic structures. Predicted domains and functional sites are readily identifiable by computer programs well known and readily available to those of skill in the art (e.g., PROSITE analysis). The results of one such analysis are provided in Figure 2.

Polypeptides often contain amino acids other than the 20 amino acids commonly referred to as the 20 naturally occurring amino acids. Further, many amino acids, including the terminal amino acids, may be modified by natural processes, such as processing and other post-translational modifications, or by chemical modification techniques well known in the art. Common modifications that occur naturally in proteins are described in basic texts, detailed monographs, and the research literature, and they are well known to those of skill in the art (some of these features are identified in Figure 2).

Known modifications include, but are not limited to, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent crosslinks, formation of cystine, formation of pyroglutamate, formylation, gamma carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination.

Such modifications are well known to those of skill in the art and have been described in great detail in the scientific literature. Several particularly common modifications, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, for instance, are described in most basic texts, such as *Proteins - Structure and Molecular Properties*, 2nd Ed., T.E. Creighton, W. H. Freeman and Company, New York (1993). Many detailed reviews are available on this subject, such as by Wold, F., *Posttranslational Covalent Modification of*

*Proteins*, B.C. Johnson, Ed., Academic Press, New York 1-12 (1983); Seifter *et al.* (*Meth. Enzymol.* 182: 626-646 (1990)) and Rattan *et al.* (*Ann. N.Y. Acad. Sci.* 663:48-62 (1992)).

Accordingly, the proteins of the present invention also encompass derivatives or analogs in which a substituted amino acid residue is not one encoded by the genetic code, in which a substituent group is included, in which the mature protein of the present invention is fused with another compound, such as a compound to increase the half-life of the protein of the present invention (for example, polyethylene glycol), or in which the additional amino acids are fused to the mature protein of the present invention, such as a leader or secretory sequence or a sequence for purification of the mature protein of the present invention or a pro-protein sequence.

#### Protein/Peptide Uses

The proteins of the present invention can be used in substantial and specific assays related to the functional information provided in the Figures: to label agents with tagged sulfur, to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its binding partner or ligand) in biological fluids; to metabolize compounds for toxicity studies; and as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state). Where the protein binds or potentially binds to another protein or ligand (such as, for example, in a enzyme-effector protein interaction or enzyme-ligand interaction), the protein can be used to identify the binding partner/ligand so as to develop a system to identify inhibitors of the binding interaction. Any or all of these uses are capable of being developed into reagent grade or kit format for commercialization as commercial products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E. F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S. L. and A. R. Kimmel eds., 1987.

The potential uses of the proteins of the present invention are based primarily on the source of the protein as well as the class/action of the protein. For example, proteins related to members of the sulfotransferase proteins are typically drug-metabolizing enzymes and serve as targets for identifying agents for use in mammalian therapeutic applications, e.g. a human drug, particularly in modulating a biological or pathological response in a cell or tissue that expresses the protein of the present invention. Experimental data provided in Figure 1 shows that the proteins of the present



invention are expressed at least in the kidney and small intestine. Specifically, multiple cDNA libraries were screened and positive clones were retrieved from libraries generated from mRNA isolated from these tissues. A large percentage of pharmaceutical agents are being developed that modulate the activity of proteins, particularly members of the sulfotransferase subfamily (see  
5 Background of the Invention) and proteins that are expressed in the kidney and intestine (killing cancer cells of these tissues that express these proteins). The structural and functional information provided in the Background and Figures provide specific and substantial uses for the molecules of the present invention, particularly in combination with the expression information provided in Figure 1. Experimental data provided in Figure 1 shows that the sulfotransferase of the present  
10 invention is expressed at least in the kidney and small intestine. Further, the proteins of the present invention can be used in an industrial setting to label agents with tagged sulfur. Such uses can readily be determined using the information provided herein, that which is known in the art, and routine experimentation.

The proteins of the present invention (including variants and fragments that may have been  
15 disclosed prior to the present invention) are useful for biological assays related to proteins that are related to members of the sulfotransferase subfamily. Such assays involve any of the known drug-metabolizing enzyme functions or activities or properties useful for diagnosis and treatment of drug-metabolizing enzyme-related conditions that are specific for the subfamily of drug-metabolizing enzymes that the one of the present invention belongs to, particularly in cells and tissues that express  
20 the drug-metabolizing enzyme. Experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine. Specifically, multiple cDNA libraries were screened and positive clones were retrieved from libraries generated from mRNA isolated from these tissues.

The proteins of the present invention are also useful in labeling target agents with tagged sulfur.  
25 Such uses are readily apparent based on the enzymatic properties of the proteins of the present invention.

The proteins of the present invention are also useful in drug screening assays, in cell-based or cell-free systems. Cell-based systems can be native, i.e., cells that normally express the protein of the present invention, as a biopsy or expanded in cell culture. Experimental data provided in Figure 1  
30 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine. In an alternate embodiment, cell-based assays involve recombinant host cells expressing the protein of the present invention.

The polypeptides can be used to identify compounds that modulate the activity of the protein in its natural state or an altered form that causes a specific disease or pathology associated with the

protein. Both the protein of the present invention and appropriate variants and fragments can be used in high-throughput screens to assay candidate compounds for the ability to bind to the protein of the present invention. These compounds can be further screened against a functional protein of the present invention to determine the effect of the compound on the protein's activity. Further, these compounds can be tested in animal or invertebrate systems to determine activity/effectiveness. Compounds can be identified that activate (agonist) or inactivate (antagonist) the protein of the present invention to a desired degree. Further, antagonists can be used to quench chemical reactions of the protein (e.g. a labeling step).

Further, the proteins of the present invention can be used to screen a compound for the ability to stimulate or inhibit interaction between the protein and a molecule that normally interacts with the protein. Such assays typically include the steps of combining the protein of the present invention with a candidate compound under conditions that allow the protein of the present invention, or fragment, to interact with the target molecule, and to detect the formation of a complex between the protein and the target or to detect the biochemical consequence of the interaction with the protein of the present invention and the target.

Candidate compounds include, for example, 1) peptides such as soluble peptides, including Ig-tailed fusion peptides and members of random peptide libraries (see, e.g., Lam *et al.*, *Nature* 354:82-84 (1991); Houghten *et al.*, *Nature* 354:84-86 (1991)) and combinatorial chemistry-derived molecular libraries made of D- and/or L- configuration amino acids; 2) phosphopeptides (e.g., members of random and partially degenerate, directed phosphopeptide libraries, see, e.g., Songyang *et al.*, *Cell* 72:767-778 (1993)); 3) antibodies (e.g., polyclonal, monoclonal, humanized, anti-idiotypic, chimeric, and single chain antibodies as well as Fab, F(ab')<sub>2</sub>, Fab expression library fragments, and epitope-binding fragments of antibodies); and 4) small organic and inorganic molecules (e.g., molecules obtained from combinatorial and natural product libraries).

One candidate compound is a soluble fragment of the protein that competes for substrate binding. Other candidate compounds include mutant proteins or appropriate fragments containing mutations that affect protein function and thus compete for substrate. Accordingly, a fragment that competes for substrate, for example with a higher affinity, or a fragment that binds substrate but does not allow release, is encompassed by the invention.

Any of the biological or biochemical functions mediated by the protein of the present invention can be used as an endpoint assay. These include all of the biochemical or biochemical/biological events described herein, in the references cited herein, incorporated by reference for these endpoint assay targets, and other functions known to those of ordinary skill in the art or that can be readily identified using the information provided in the Figures, particularly Figure 2. Specifically, a biological

function of a cell or tissues that expresses the protein of the present invention can be assayed. Experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine. Specifically, multiple cDNA libraries were screened and positive clones were retrieved from libraries generated from mRNA isolated from these tissues.

Binding and/or activating compounds can also be screened by using chimeric protein of the present invention in which the amino terminal extracellular domain, or parts thereof, the entire transmembrane domain or subregions, such as any of the seven transmembrane segments or any of the intracellular or extracellular loops and the carboxy terminal intracellular domain, or parts thereof, can be replaced by heterologous domains or subregions. For example, a substrate-binding region can be used that interacts with a different substrate than that which is recognized by the native protein of the present invention. Accordingly, a different set of signal transduction components is available as an end-point assay for activation. This allows for assays to be performed in other than the specific host cell from which the protein of the present invention is derived.

The proteins of the present invention are also useful in competition binding assays in methods designed to discover compounds that interact with the protein of the present invention (e.g. binding partners and/or ligands). Thus, a compound is exposed to a protein of the present invention under conditions that allow the compound to bind or to otherwise interact with the polypeptide. Soluble protein of the present invention polypeptide is also added to the mixture. If the test compound interacts with the soluble protein of the present invention polypeptide, it decreases the amount of complex formed or activity from the protein of the present invention target. This type of assay is particularly useful in cases in which compounds are sought that interact with specific regions of the protein of the present invention. Thus, the soluble polypeptide that competes with the target protein of the present invention region is designed to contain peptide sequences corresponding to the region of interest.

To perform cell free drug screening assays, it is sometimes desirable to immobilize either the protein of the present invention, or fragment, or its target molecule to facilitate separation of complexes from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay.

Techniques for immobilizing proteins on matrices can be used in the drug screening assays. In one embodiment, a fusion protein can be provided which adds a domain that allows the protein to be bound to a matrix. For example, glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the cell lysates (e.g., <sup>35</sup>S-labeled) and the candidate compound, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological

conditions for salt and pH). Following incubation, the beads are washed to remove any unbound label, and the matrix immobilized and radiolabel determined directly, or in the supernatant after the complexes are dissociated. Alternatively, the complexes can be dissociated from the matrix, separated by SDS-PAGE, and the level of protein of the present invention-binding protein found in the bead  
5 fraction quantitated from the gel using standard electrophoretic techniques. For example, either the polypeptide or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin using techniques well known in the art. Alternatively, antibodies reactive with the protein but which do not interfere with binding of the protein to its target molecule can be derivatized to the wells of the plate, and the protein trapped in the wells by antibody conjugation. Preparations of a protein of the  
10 present invention-binding protein and a candidate compound are incubated in the protein of the present invention-presenting wells and the amount of complex trapped in the well can be quantitated. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the protein of the present invention target molecule, or which are reactive with protein of the present invention and  
15 compete with the target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the target molecule.

Agents that modulate one of the proteins of the present invention can be identified using one or more of the above assays, alone or in combination. It is generally preferable to use a cell-based or cell free system first and then confirm activity in an animal or other model system. Such model systems  
20 are well known in the art and can readily be employed in this context.

Modulators of protein of the present invention activity identified according to these drug screening assays can be used to treat a subject with a disorder mediated by the protein of the present invention pathway, by treating cells or tissues that express the protein of the present invention. Experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is  
25 expressed at least in the kidney and small intestine. These methods of treatment include the steps of administering a modulator of protein of the present invention activity in a pharmaceutical composition to a subject in need of such treatment, the modulator being identified as described herein.

In yet another aspect of the invention, the protein of the present inventions can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, e.g., U.S. Patent No. 5,283,317; Zervos *et al.* (1993) *Cell* 72:223-232; Madura *et al.* (1993) *J. Biol. Chem.* 268:12046-12054; Bartel  
30 *et al.* (1993) *Biotechniques* 14:920-924; Iwabuchi *et al.* (1993) *Oncogene* 8:1693-1696; and Brent WO94/10300), to identify other proteins, which bind to or interact with the protein of the present invention and are involved in the protein's enzymatic activity.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for a protein of the present invention is fused to a gene encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a protein of the present invention-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein which interacts with the protein of the present invention.

This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an agent identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a protein of the present invention-modulating agent, an antisense protein of the present invention nucleic acid molecule, a protein of the present invention-specific antibody, or a protein of the present invention-binding partner) can be used in an animal or other model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal or other model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above-described screening assays for treatments as described herein.

The proteins of the present invention are also useful to provide a target for diagnosing a disease or predisposition to disease mediated by the peptide. Accordingly, the invention provides methods for detecting the presence, or levels of, the protein (or encoding mRNA) in a cell, tissue, or organism. Experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine. The method involves contacting a biological sample with a compound capable of interacting with the protein of the present invention such that the interaction can be detected. Such an assay can be provided in a single detection format or a multi-detection format such as an antibody chip array.

One agent for detecting a protein in a sample is an antibody capable of selectively binding to protein. A biological sample includes tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject.

The proteins of the present invention also provide targets for diagnosing active protein activity, disease, or predisposition to disease, in a patient having a variant peptide, particularly activities and conditions that are known for other members of the family of proteins to which the present one belongs. Thus, the peptide can be isolated from a biological sample and assayed for the presence of a genetic mutation that results in aberrant peptide. This includes amino acid substitution, deletion, insertion, rearrangement, (as the result of aberrant splicing events), and inappropriate post-translational modification. Analytic methods include altered electrophoretic mobility, altered tryptic peptide digest, altered protein of the present invention activity in cell-based or cell-free assay, alteration in substrate or antibody-binding pattern, altered isoelectric point, direct amino acid sequencing, and any other of the known assay techniques useful for detecting mutations in a protein. Such an assay can be provided in a single detection format or a multi-detection format such as an antibody chip array.

*In vitro* techniques for detection of peptide include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence using a detection reagent, such as an antibody or protein binding agent. Alternatively, the peptide can be detected *in vivo* in a subject by introducing into the subject a labeled anti-peptide antibody or other types of detection agent. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques. Particularly useful are methods that detect the allelic variant of a peptide expressed in a subject and methods which detect fragments of a peptide in a sample.

The peptides are also useful in pharmacogenomic analysis. Pharmacogenomics deal with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, e.g., Eichelbaum, M. (*Clin. Exp. Pharmacol. Physiol.* 23(10-11):983-985 (1996)), and Linder, M.W. (*Clin. Chem.* 43(2):254-266 (1997)). The clinical outcomes of these variations result in severe toxicity of therapeutic drugs in certain individuals or therapeutic failure of drugs in certain individuals as a result of individual variation in metabolism. Thus, the genotype of the individual can determine the way a therapeutic compound acts on the body or the way the body metabolizes the compound. Further, the activity of drug metabolizing enzymes effects both the intensity and duration of drug action. Thus, the pharmacogenomics of the individual permit the selection of effective compounds and effective dosages of such compounds for prophylactic or therapeutic treatment based on the individual's genotype. The discovery of genetic polymorphisms in some drug metabolizing enzymes has explained why some patients do not obtain the expected drug effects, show an exaggerated drug effect, or experience serious toxicity from standard drug dosages. Polymorphisms can be expressed in the phenotype of the extensive metabolizer and the phenotype of the poor metabolizer. Accordingly, genetic polymorphism may lead to allelic protein variants of the

protein of the present invention in which one or more of the protein of the present invention functions in one population is different from those in another population. The peptides thus allow a target to ascertain a genetic predisposition that can affect treatment modality. Thus, in a ligand-based treatment, polymorphism may give rise to amino terminal extracellular domains and/or other substrate-binding regions that are more or less active in substrate binding. and protein of the present invention activation. Accordingly, substrate dosage would necessarily be modified to maximize the therapeutic effect within a given population containing a polymorphism. As an alternative to genotyping, specific polymorphic peptides could be identified.

The peptides are also useful for treating a disorder characterized by an absence of, inappropriate, or unwanted expression of the protein. Experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine. Accordingly, methods for treatment include the use of the protein of the present invention or fragments.

#### Antibodies

The invention also provides antibodies that selectively bind to one of the proteins of the present invention, a protein comprising a fragment of such a protein, as well as variants and fragments thereof. As used herein, an antibody selectively binds a target protein or peptide when it binds the target peptide and does not significantly bind to unrelated peptides or proteins. An antibody is still considered to selectively bind a peptide even if it also binds to other proteins that are not substantially homologous with the target peptide so long as such proteins share homology with a fragment or domain of the peptide target of the antibody. In this case, it would be understood that antibody binding to the peptide is still selective despite some degree of cross-reactivity.

As used herein, an antibody is defined in terms consistent with that recognized within the art: they are multi-subunit proteins produced by a mammalian organism in response to an antigen challenge. The antibodies of the present invention include polyclonal antibodies and monoclonal antibodies, as well as fragments of such antibodies, including, but not limited to, Fab or F(ab')<sub>2</sub>, and Fv fragments.

Many methods are known for generating and/or identifying antibodies to a given target peptide. Several such methods are described by Harlow, Antibodies, Cold Spring Harbor Press, (1989).

In general, to generate antibodies, an isolated peptide is used as an immunogen and is administered to a mammalian organism, such as a rat, rabbit or mouse. The full-length protein, an antigenic peptide fragment or a fusion protein can be used. Particularly important fragments are those covering functional domains, such as the domains identified in Figure 2, and domain of sequence

homology or divergence amongst the family, such as those that can readily be identified using protein alignment methods and as presented in the Figures.

Antibodies are preferably prepared from regions or discrete fragments of the protein of the present invention. Antibodies can be prepared from any region of the peptide as described herein.

5 However, preferred regions will include those involved in function/activity and/or protein/binding partner interaction. Figure 2 can be used to identify particularly important regions while sequence alignment can be used to identify conserved and unique sequence fragments.

An antigenic fragment will typically comprise at least 8 contiguous amino acid residues. The antigenic peptide can comprise, however, at least 10, 12, 14, 16 or more amino acid residues. Such  
10 fragments can be selected on a physical property, such as fragments correspond to regions that are located on the surface of the protein, e.g., hydrophilic regions or can be selected based on sequence uniqueness (see Figure 2).

Detection on an antibody of the present invention can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance. Examples of detectable substances include various  
15 enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase,  $\beta$ -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include  
20 umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{35}\text{S}$  or  $^3\text{H}$ .

#### Antibody Uses

25 The antibodies can be used to isolate one of the proteins of the present invention by standard techniques, such as affinity chromatography or immunoprecipitation. The antibodies can facilitate the purification of the natural protein from cells and recombinantly produced protein expressed in host cells. In addition, such antibodies are useful to detect the presence of one of the proteins of the present invention in cells or tissues to determine the pattern of expression of the protein among various tissues  
30 in an organism and over the course of normal development. Experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine. Specifically, multiple cDNA libraries were screened and positive clones were retrieved from libraries generated from mRNA isolated from these tissues. Further, such antibodies can be used to detect protein *in situ*, *in vitro*, or in a cell lysate or supernatant in order to evaluate the abundance and



pattern of expression. Also, such antibodies can be used to assess abnormal tissue distribution or abnormal expression during development or progression of a biological condition. Antibody detection of circulating fragments of the full-length protein can be used to identify protein turnover.

Further, the antibodies can be used to assess expression in disease states such as in active stages of the disease or in an individual with a predisposition toward disease related to the protein's function. When a disorder is caused by an inappropriate tissue distribution, developmental expression, level of expression of the protein, or expressed/processed form, the antibody can be prepared against the normal protein. Experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine. If a disorder is characterized by a specific mutation in the protein, antibodies specific for this mutant protein can be used to assay for the presence of the specific mutant protein.

The antibodies can also be used to assess normal and aberrant subcellular localization of cells in the various tissues in an organism. Experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine. The diagnostic uses can be applied, not only in genetic testing, but also in monitoring a treatment modality. Accordingly, where treatment is ultimately aimed at correcting expression level or the presence of aberrant sequence and aberrant tissue distribution or developmental expression, antibodies directed against the protein or relevant fragments can be used to monitor therapeutic efficacy.

Additionally, antibodies are useful in pharmacogenomic analysis. Thus, antibodies prepared against polymorphic proteins can be used to identify individuals that require modified treatment modalities. The antibodies are also useful as diagnostic tools as an immunological marker for aberrant protein analyzed by electrophoretic mobility, isoelectric point, tryptic peptide digest, and other physical assays known to those in the art.

The antibodies are also useful for tissue typing. Experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine. Thus, where a specific protein has been correlated with expression in a specific tissue, antibodies that are specific for this protein can be used to identify a tissue type.

The antibodies are also useful for inhibiting protein function, for example, blocking the binding of the protein to a binding partner such as a substrate. These uses can also be applied in a therapeutic context in which treatment involves inhibiting the protein's function. An antibody can be used, for example, to block binding, thus modulating (agonizing or antagonizing) the protein's activity. Antibodies can be prepared against specific fragments containing sites required for function or against intact protein that is associated with a cell or cell membrane. See Figure 2 for structural information relating to the proteins of the present invention.

The invention also encompasses kits for using antibodies to detect the presence of a protein in a biological sample. The kit can comprise antibodies such as a labeled or labelable antibody and a compound or agent for detecting protein in a biological sample; means for determining the amount of protein in the sample; means for comparing the amount of protein in the sample with a standard; and instructions for use. Such a kit can be supplied to detect a single protein or epitope or can be configured to detect one of a multitude of epitopes, such as in an antibody detection array. Arrays are described in detail below for nucleic acids but similar methods have been developed for antibody arrays,

#### Nucleic Acid Molecules

The present invention further provides isolated nucleic acid molecules that encode a protein of the present invention (cDNA, transcript and genomic sequence). Such nucleic acid molecules will consist of, consist essentially of, or comprise a nucleotide sequence that encodes one of the proteins of the present invention, an allelic variant thereof, or an ortholog or paralog thereof.

As used herein, an "isolated" nucleic acid molecule is one that is separated from other nucleic acid present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. However, there can be some flanking nucleotide sequences, for example up to about 5KB, 4KB, 3KB, 2KB, or 1KB or less, particularly contiguous peptide encoding sequences and peptide encoding sequences within the same gene but separated by introns in the genomic sequence. The important point is that the nucleic acid is isolated from remote and unimportant flanking sequences such that it can be subjected to the specific manipulations described herein such as recombinant expression, preparation of probes and primers, and other uses specific to the nucleic acid sequences.

Moreover, an "isolated" nucleic acid molecule, such as a transcript/cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when chemically synthesized. However, the nucleic acid molecule can be fused to other coding or regulatory sequences and still be considered isolated.

For example, recombinant DNA molecules contained in a vector are considered isolated. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the isolated DNA molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced synthetically.

Accordingly, the present invention provides nucleic acid molecules that consist of the nucleotide sequence shown in Figure 1 or 3 (SEQ ID NOS:1 and 2, alternative transcript sequences, and SEQ ID NO:3, cDNA sequences, transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NOS:4-6. A nucleic acid molecule consists of a nucleotide sequence when the nucleotide sequence is the complete nucleotide sequence of the nucleic acid molecule.

The present invention further provides nucleic acid molecules that consist essentially of the nucleotide sequence shown in Figure 1 or 3 (SEQ ID NOS:1 and 2, alternative transcript sequences, and SEQ ID NO:3, cDNA sequences, transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NOS:4-6. A nucleic acid molecule consists essentially of a nucleotide sequence when such a nucleotide sequence is present with only a few additional nucleic acid residues in the final nucleic acid molecule.

The present invention further provides nucleic acid molecules that comprise the nucleotide sequences shown in Figure 1 or 3 (SEQ ID NOS:1 and 2, alternative transcript sequences, and SEQ ID NO:3, cDNA sequences, transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NOS:4-6. A nucleic acid molecule comprises a nucleotide sequence when the nucleotide sequence is at least part of the final nucleotide sequence of the nucleic acid molecule. In such a fashion, the nucleic acid molecule can be only the nucleotide sequence or have additional nucleic acid residues, such as nucleic acid residues that are naturally associated with it or heterologous nucleotide sequences. Such a nucleic acid molecule can have a few additional nucleotides or can comprises several hundred or more additional nucleotides. A brief description of how various types of these nucleic acid molecules can be readily made/isolated is provided below.

In Figures 1 and 3, both coding and non-coding sequences are provided. Because of the source of the present invention, humans genomic sequence (Figure 3) and cDNA/transcript sequences (Figure 1), the nucleic acid molecules in the Figures will contain genomic intronic sequences, 5' and 3' non-coding sequences, gene regulatory regions and non-coding intergenic sequences. In general such sequence features are either noted in Figures 1 and 3 or can readily be identified using computational tools known in the art. As discussed below, some of the non-coding regions, particularly gene regulatory elements such as promoters, are useful for a variety of purposes, e.g. control of heterologous gene expression, target for identifying gene activity modulating compounds, and are particularly claimed as fragments of the genomic sequence provided herein.

The isolated nucleic acid molecules can encode the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature peptide (when the mature form has more than one peptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, facilitate protein trafficking, prolong or shorten protein half-life or facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in situ*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

As mentioned above, the isolated nucleic acid molecules include, but are not limited to, the sequence encoding the proteins of the present invention alone, the sequence encoding the mature peptide and additional coding sequences, such as a leader or secretory sequence (e.g., a pre-pro or pro-protein sequence), the sequence encoding the mature peptide, with or without the additional coding sequences, plus additional non-coding sequences, for example introns and non-coding 5' and 3' sequences such as transcribed but non-translated sequences that play a role in transcription, mRNA processing (including splicing and polyadenylation signals), ribosome binding and stability of mRNA. In addition, the nucleic acid molecule may be fused to a marker sequence encoding, for example, a peptide that facilitates purification.

Isolated nucleic acid molecules can be in the form of RNA, such as mRNA, or in the form of DNA, including cDNA and genomic DNA obtained by cloning or produced by chemical synthetic techniques or by a combination thereof. The nucleic acid, especially DNA, can be double-stranded or single-stranded. Single-stranded nucleic acid can be the coding strand (sense strand) or the non-coding strand (anti-sense strand).

The invention further provides nucleic acid molecules that encode fragments of the proteins of the present invention as well as nucleic acid molecules that encode obvious variants of the proteins of the present invention that are described above. Such nucleic acid molecules may be naturally occurring, such as allelic variants (same locus), paralogs (different locus), and orthologs (different organism), or may be constructed by recombinant DNA methods or by chemical synthesis. Such non-naturally occurring variants may be made by mutagenesis techniques, including those applied to nucleic acid molecules, cells, or organisms. Accordingly, as discussed above, the variants can contain nucleotide substitutions, deletions, inversions and insertions. Variation can occur in either or both the coding and non-coding regions. The variations can produce both conservative and non-conservative amino acid substitutions.

The present invention further provides non-coding fragments of the nucleic acid molecules provided in Figures 1 and 3. Preferred non-coding fragments include, but are not limited to, promoter sequences, enhancer sequences, gene modulating sequences and gene termination sequences. Such

fragments are useful in controlling heterologous gene expression and in developing screens to identify gene-modulating agents. A promoter can readily be identified as being 5' to the ATG start site in the genomic sequence provided in Figure 3.

A fragment comprises a contiguous nucleotide sequence greater than 12 or more nucleotides. Further, a fragment could at least 30, 40, 50, 100, 250 or 500 nucleotides in length. The length of the fragment will be based on its intended use. For example, the fragment can encode epitope bearing regions of the peptide, or can be useful as DNA probes and primers. Such fragments can be isolated using the known nucleotide sequence to synthesize an oligonucleotide probe. A labeled probe can then be used to screen a cDNA library, genomic DNA library, or mRNA to isolate nucleic acid corresponding to the coding region. Further, primers can be used in PCR reactions to clone specific regions of gene.

A probe/primer typically comprises substantially a purified oligonucleotide or oligonucleotide pair. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, 20, 25, 40, 50 or more consecutive nucleotides.

Orthologs, homologs, and allelic variants can be identified using methods well known in the art. As described in the Peptide Section, these variants comprise a nucleotide sequence encoding a peptide that is typically 60-70%, 70-80%, 80-90%, and more typically at least about 90-95% or more homologous to the nucleotide sequence shown in the Figure sheets or a fragment of this sequence. Such nucleic acid molecules can readily be identified as being able to hybridize under moderate to stringent conditions, to the nucleotide sequence shown in the Figure sheets or a fragment of the sequence. Allelic variants can readily be determined by genetic locus of the encoding gene. As indicated by the data presented in Figure 3, the map position was determined to be on chromosome 2 near marker SHGC-1639 (LOD score 10.01) using radiation hybrid mapping.

Figure 3 provides SNP information that was identified from chromosomal sequence obtained from a number of individuals as well as SNPs found during the cDNA cloning process. These include, a6094g, t7898c, a10018c, a10791t, a12451t, t14674c, g14860c, a20133g, a1025c, t12602c, g12829t, a1442t.

As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences encoding a peptide at least 60-70% homologous to each other typically remain hybridized to each other. The conditions can be such that sequences at least about 60%, at least about 70%, or at least about 80% or more homologous to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. One example of stringent hybridization conditions are hybridization in 6X

sodium chloride/sodium citrate (SSC) at about 45C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65C. Examples of moderate to low stringency hybridization conditions are well known in the art.

5        Nucleic Acid Molecule Uses

The nucleic acid molecules of the present invention are useful for probes, primers, chemical intermediates, mapping reagents, expression control reagents and in biological assays. The nucleic acid molecules are useful as a hybridization probe for messenger RNA, transcript/cDNA and genomic DNA to isolate full-length cDNA and genomic clones encoding the peptide described in Figure 2 and  
10 to isolate cDNA and genomic clones that correspond to variants (alleles, orthologs, etc.) producing the same or related peptides shown in Figure 2. As illustrated in Figure 3, identified SNP variations include a1008c, a6094g, t7898c, a10018c, a10791t, a12451t, t14674c, g14860c, a20133g, a1025c, t12602c, g12829t, a1442t.

The probe can correspond to any sequence along the entire length of the nucleic acid molecules  
15 provided in the Figures. Accordingly, it could be derived from 5' noncoding regions, the coding region, and 3' noncoding regions. However, as discussed, fragments are not to be construed as encompassing fragments disclosed prior to the present invention.

The nucleic acid molecules are also useful as primers for PCR to amplify any given region of a nucleic acid molecule and are useful to synthesize antisense molecules of desired length and sequence.  
20 Such molecules are useful in genotyping and mapping experiments, particularly if the probed region contains one or more of the sequence variants described herein.

The nucleic acid molecules are also useful for constructing recombinant vectors. Such vectors include expression vectors that express a portion of, or all of, the peptide sequences. Vectors also include insertion vectors, used to integrate into another nucleic acid molecule sequence, such as into  
25 the cellular genome, to alter *in situ* expression of a gene and/or gene product. For example, an endogenous coding sequence can be replaced via homologous recombination with all or part of the coding region containing one or more specifically introduced mutations.

The nucleic acid molecules are also useful for expressing antigenic portions of the proteins.

The nucleic acid molecules are also useful as probes for determining the chromosomal  
30 positions of the nucleic acid molecules by means of *in situ* hybridization methods. As indicated by the data presented in Figure 3, the map position was determined to be on chromosome 2 near marker SHGC-1639 (LOD score 10.01) using radiation hybrid mapping. Further, the SNP variants provided herein can be used in mapping and linkage studies.

The nucleic acid molecules are also useful in making vectors containing the gene regulatory regions of the nucleic acid molecules of the present invention.

The nucleic acid molecules are also useful for designing ribozymes corresponding to all, or a part, of the mRNA produced from the nucleic acid molecules described herein.

5 The nucleic acid molecules are also useful for making vectors that express part, or all, of the peptides.

The nucleic acid molecules are also useful for constructing host cells expressing a part, or all, of the nucleic acid molecules and peptides.

10 The nucleic acid molecules are also useful for constructing transgenic animals expressing all, or a part, of the nucleic acid molecules and peptides.

The nucleic acid molecules are also useful as hybridization probes for determining the presence, level, form and distribution of nucleic acid expression. Experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine. Specifically, multiple cDNA libraries were screened and positive clones were retrieved from  
15 libraries generated from mRNA isolated from these tissues. Accordingly, the probes can be used to detect the presence of, or to determine levels of, a specific nucleic acid molecule in cells, tissues, and in organisms. The nucleic acid whose level is determined can be DNA or RNA. Accordingly, probes corresponding to the peptides described herein can be used to assess expression and/or gene copy number in a given cell, tissue, or organism. These uses are relevant for diagnosis of disorders involving  
20 an increase or decrease in expression relative to normal results.

*In vitro* techniques for detection of mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detecting DNA includes Southern hybridizations and *in situ* hybridization.

Probes can be used as a part of a diagnostic test kit for identifying cells or tissues that express a  
25 protein of the present invention, such as by measuring a level of a protein of the present invention-encoding nucleic acid in a sample of cells from a subject e.g., mRNA or genomic DNA, or determining if a protein of the present invention gene has been mutated. Experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine. Specifically, multiple cDNA libraries were screened and positive clones were retrieved from  
30 libraries generated from mRNA isolated from these tissues.

Nucleic acid expression assays are useful for drug screening to identify compounds that modulate protein of the present invention nucleic acid expression.

The invention thus provides a method for identifying a compound that can be used to treat a disorder associated with nucleic acid expression of the protein of the present invention gene,

particularly biological and pathological processes that are mediated by the protein of the present invention in cells and tissues that express it. Experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine. The method typically includes assaying the ability of the compound to modulate the expression of the protein of the present invention nucleic acid and thus identifying a compound that can be used to treat a disorder characterized by undesired protein of the present invention nucleic acid expression. The assays can be performed in cell-based and cell-free systems. Cell-based assays include cells naturally expressing the protein of the present invention nucleic acid or recombinant cells genetically engineered to express specific nucleic acid sequences.

Thus, modulators of protein of the present invention gene expression can be identified in a method wherein a cell is contacted with a candidate compound and the expression of mRNA determined. The level of expression of protein of the present invention mRNA in the presence of the candidate compound is compared to the level of expression of protein of the present invention mRNA in the absence of the candidate compound. The candidate compound can then be identified as a modulator of nucleic acid expression based on this comparison and be used, for example to treat a disorder characterized by aberrant nucleic acid expression. When expression of mRNA is statistically significantly greater in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of nucleic acid expression. When nucleic acid expression is statistically significantly less in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of nucleic acid expression.

The invention further provides methods of treatment, with the nucleic acid as a target, using a compound identified through drug screening as a gene modulator to modulate protein of the present invention nucleic acid expression in cells and tissues that express the protein. Experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine. Specifically, multiple cDNA libraries were screened and positive clones were retrieved from libraries generated from mRNA isolated from these tissues. Modulation includes both up-regulation (i.e. activation or agonization) or down-regulation (suppression or antagonization) or nucleic acid expression.

Alternatively, a modulator for nucleic acid expression can be a small molecule or drug identified using the screening assays described herein as long as the drug or small molecule inhibits the protein of the present invention nucleic acid expression in the cells and tissues that express the protein. Experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine.



The nucleic acid molecules are also useful for monitoring the effectiveness of modulating compounds on the expression or activity of the protein of the present invention gene in clinical trials or in a treatment regimen. Thus, the gene expression pattern can serve as a barometer for the continuing effectiveness of treatment with the compound, particularly with compounds to which a patient can develop resistance. The gene expression pattern can also serve as a marker indicative of a physiological response of the affected cells to the compound. Accordingly, such monitoring would allow either increased administration of the compound or the administration of alternative compounds to which the patient has not become resistant. Similarly, if the level of nucleic acid expression falls below a desirable level, administration of the compound could be commensurately decreased.

The nucleic acid molecules are also useful in diagnostic assays for qualitative changes in protein of the present invention nucleic acid expression, and particularly in qualitative changes that lead to pathology. The nucleic acid molecules can be used to detect mutations in protein of the present invention genes and gene expression products such as mRNA. The nucleic acid molecules can be used as hybridization probes to detect naturally occurring genetic mutations in the protein of the present invention gene and thereby to determine whether a subject with the mutation is at risk for a disorder caused by the mutation. Mutations include deletion, addition, or substitution of one or more nucleotides in the gene, chromosomal rearrangement, such as inversion or transposition, modification of genomic DNA, such as aberrant methylation patterns or changes in gene copy number, such as amplification. Detection of a mutated form of the protein of the present invention gene associated with a dysfunction provides a diagnostic tool for an active disease or susceptibility to disease when the disease results from overexpression, underexpression, or altered expression of a protein of the present invention.

Individuals carrying mutations in the protein of the present invention gene can be detected at the nucleic acid level by a variety of techniques. Figure 3 provides SNP information that was identified from chromosomal sequence obtained from a number of individuals as well as SNPs found during the cDNA cloning process. These include, a6094g, t7898c, a10018c, a10791t, a12451t, t14674c, g14860c, a20133g, a1025c, t12602c, g12829t, a1442t. As indicated by the data presented in Figure 3, the map position was determined to be on chromosome 2 near marker SHGC-1639 (LOD score 10.01) using radiation hybrid mapping. Genomic DNA can be analyzed directly or can be amplified by using PCR prior to analysis. RNA or cDNA can be used in the same way. In some uses, detection of the mutation involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, e.g. U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, e.g., Landegran *et al.*, *Science* 241:1077-1080 (1988); and Nakazawa *et al.*, *PNAS* 91:360-364 (1994)), the latter of which can be particularly useful for detecting point mutations in the

gene (see Abravaya *et al.*, *Nucleic Acids Res.* 23:675-682 (1995)). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (e.g., genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a gene under conditions such that hybridization and amplification of the gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. Deletions and insertions can be detected by a change in size of the amplified product compared to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to normal RNA or antisense DNA sequences.

Alternatively, mutations in a protein of the present invention gene can be directly identified, for example, by alterations in restriction enzyme digestion patterns determined by gel electrophoresis.

Further, sequence-specific ribozymes (U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site. Perfectly matched sequences can be distinguished from mismatched sequences by nuclease cleavage digestion assays or by differences in melting temperature.

Sequence changes at specific locations can also be assessed by nuclease protection assays such as RNase and S1 protection or the chemical cleavage method. Furthermore, sequence differences between a mutant protein of the present invention gene and a wild-type gene can be determined by direct DNA sequencing. A variety of automated sequencing procedures can be utilized when performing the diagnostic assays (Naeve, C.W., (1995) *Biotechniques* 19:448), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen *et al.*, *Adv. Chromatogr.* 36:127-162 (1996); and Griffin *et al.*, *Appl. Biochem. Biotechnol.* 38:147-159 (1993)).

Other methods for detecting mutations in the gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA duplexes (Myers *et al.*, *Science* 230:1242 (1985)); Cotton *et al.*, *PNAS* 85:4397 (1988); Saleeba *et al.*, *Meth. Enzymol.* 217:286-295 (1992)), electrophoretic mobility of mutant and wild type nucleic acid is compared (Orita *et al.*, *PNAS* 86:2766 (1989); Cotton *et al.*, *Mutat. Res.* 285:125-144 (1993); and Hayashi *et al.*, *Genet. Anal. Tech. Appl.* 9:73-79 (1992)), and movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (Myers *et al.*, *Nature* 313:495 (1985)). Examples of other techniques for detecting point mutations include selective oligonucleotide hybridization, selective amplification, and selective primer extension.

The nucleic acid molecules are also useful for testing an individual for a genotype that while not necessarily causing the disease, nevertheless affects the treatment modality. Thus, the nucleic acid molecules can be used to study the relationship between an individual's genotype and the individual's

response to a compound used for treatment (pharmacogenomic relationship). Accordingly, the nucleic acid molecules described herein can be used to assess the mutation content of the protein of the present invention gene in an individual in order to select an appropriate compound or dosage regimen for treatment. Figure 3 provides SNP information that was identified from chromosomal sequence  
5 obtained from a number of individuals as well as SNPs found during the cDNA cloning process. These include, a6094g, t7898c, a10018c, a10791t, a12451t, t14674c, g14860c, a20133g, a1025c, t12602c, g12829t, a1442t.

Thus nucleic acid molecules displaying genetic variations that affect treatment provide a diagnostic target that can be used to tailor treatment in an individual. Accordingly, the production of  
10 recombinant cells and animals containing these polymorphisms allow effective clinical design of treatment compounds and dosage regimens.

The nucleic acid molecules are thus useful as antisense constructs to control protein of the present invention gene expression in cells, tissues, and organisms. A DNA antisense nucleic acid molecule is designed to be complementary to a region of the gene involved in transcription, preventing  
15 transcription and hence production of protein of the present invention. An antisense RNA or DNA nucleic acid molecule would hybridize to the mRNA and thus block translation of mRNA into a protein of the present invention.

Alternatively, a class of antisense molecules can be used to inactivate mRNA in order to decrease expression of protein of the present invention nucleic acid. Accordingly, these molecules can  
20 treat a disorder characterized by abnormal or undesired protein of the present invention nucleic acid expression. This technique involves cleavage by means of ribozymes containing nucleotide sequences complementary to one or more regions in the mRNA that attenuate the ability of the mRNA to be translated. Possible regions include coding regions and particularly coding regions corresponding to the catalytic and other functional activities of the protein of the present invention, such as substrate  
25 binding.

The nucleic acid molecules also provide vectors for gene therapy in patients containing cells that are aberrant in gene expression. Thus, recombinant cells, which include the patient's cells that have been engineered *ex vivo* and returned to the patient, are introduced into an individual where the cells produce the desired protein of the present invention to treat the individual.

30 The invention also encompasses kits for detecting the presence of a protein of the present invention encoding nucleic acid in a biological sample. Experimental data provided in Figure 1 shows that the sulfotransferase of the present invention is expressed at least in the kidney and small intestine. Specifically, multiple cDNA libraries were screened and positive clones were retrieved from libraries generated from mRNA isolated from these tissues. For example, the kit can comprise reagents such as

a labeled or labelable nucleic acid or agent capable of detecting protein of the present invention encoding nucleic acid in a biological sample; means for determining the amount of protein of the present invention encoding nucleic acid in the sample; and means for comparing the amount of protein of the present invention encoding nucleic acid in the sample with a standard. The compound or agent  
5 can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect a protein of the present invention mRNA or DNA.

#### Nucleic Acid Arrays

The present invention further provides nucleic acid detection kits, such as arrays or  
10 microarrays of nucleic acid molecules that are based on the sequence information provided in Figures 1 and 3 (SEQ ID NOS:1 and 3).

As used herein "Arrays" or "Microarrays" refers to an array of distinct polynucleotides or oligonucleotides synthesized on a substrate, such as paper, nylon or other type of membrane, filter, chip, glass slide, or any other suitable solid support. In one embodiment, the microarray is prepared  
15 and used according to the methods described in US Patent 5,837,832, Chee *et al.*, PCT application W095/11995 (Chee *et al.*), Lockhart, D. J. *et al.* (1996; Nat. Biotech. 14: 1675-1680) and Schena, M. *et al.* (1996; Proc. Natl. Acad. Sci. 93: 10614-10619), all of which are incorporated herein in their entirety by reference. In other embodiments, such arrays are produced by the methods described by Brown *et al.*, US Patent No. 5,807,522.

The microarray or detection kit is preferably composed of a large number of unique, single-stranded nucleic acid sequences, usually either synthetic antisense oligonucleotides or fragments of cDNAs, fixed to a solid support. The oligonucleotides are preferably about 6-60 nucleotides in length, more preferably 15-30 nucleotides in length, and most preferably about 20-25 nucleotides in length. For a certain type of microarray or detection kit, it may be preferable to use oligonucleotides  
25 that are only 7-20 nucleotides in length. The microarray or detection kit may contain oligonucleotides that cover the known 5', or 3', sequence, sequential oligonucleotides which cover the full length sequence; or unique oligonucleotides selected from particular areas along the length of the sequence. Polynucleotides used in the microarray or detection kit may be oligonucleotides that are specific to a gene or genes of interest.

In order to produce oligonucleotides to a known sequence for a microarray or detection kit,  
30 the gene(s) of interest (or an ORF identified from the contigs of the present invention) is typically examined using a computer algorithm which starts at the 5' or at the 3' end of the nucleotide sequence. Typical algorithms will then identify oligomers of defined length that are unique to the gene, have a GC content within a range suitable for hybridization, and lack predicted secondary

structure that may interfere with hybridization. In certain situations it may be appropriate to use pairs of oligonucleotides on a microarray or detection kit. The "pairs" will be identical, except for one nucleotide that preferably is located in the center of the sequence. The second oligonucleotide in the pair (mismatched by one) serves as a control. The number of oligonucleotide pairs may range  
5 from two to one million. The oligomers are synthesized at designated areas on a substrate using a light-directed chemical process. The substrate may be paper, nylon or other type of membrane, filter, chip, glass slide or any other suitable solid support.

In another aspect, an oligonucleotide may be synthesized on the surface of the substrate by using a chemical coupling procedure and an ink jet application apparatus, as described in PCT  
10 application W095/251116 (Baldeschweiler *et al.*) which is incorporated herein in its entirety by reference. In another aspect, a "gridded" array analogous to a dot (or slot) blot may be used to arrange and link cDNA fragments or oligonucleotides to the surface of a substrate using a vacuum system, thermal, UV, mechanical or chemical bonding procedures. An array, such as those described above, may be produced by hand or by using available devices (slot blot or dot blot  
15 apparatus), materials (any suitable solid support), and machines (including robotic instruments), and may contain 8, 24, 96, 384, 1536, 6144 or more oligonucleotides, or any other number between two and one million which lends itself to the efficient use of commercially available instrumentation.

In order to conduct sample analysis using a microarray or detection kit, the RNA or DNA from a biological sample is made into hybridization probes. The mRNA is isolated, and cDNA is  
20 produced and used as a template to make antisense RNA (aRNA). The aRNA is amplified in the presence of fluorescent nucleotides, and labeled probes are incubated with the microarray or detection kit so that the probe sequences hybridize to complementary oligonucleotides of the microarray or detection kit. Incubation conditions are adjusted so that hybridization occurs with precise complementary matches or with various degrees of less complementarity. After removal of  
25 nonhybridized probes, a scanner is used to determine the levels and patterns of fluorescence. The scanned images are examined to determine degree of complementarity and the relative abundance of each oligonucleotide sequence on the microarray or detection kit. The biological samples may be obtained from any bodily fluids (such as blood, urine, saliva, phlegm, gastric juices, etc.), cultured cells, biopsies, or other tissue preparations. A detection system may be used to measure the  
30 absence, presence, and amount of hybridization for all of the distinct sequences simultaneously. This data may be used for large-scale correlation studies on the sequences, expression patterns, mutations, variants, or polymorphisms among samples.

Using such arrays, the present invention provides methods to identify the expression of the proteins of the present invention. In detail, such methods comprise incubating a test sample with

one or more nucleic acid molecules and assaying for binding of the nucleic acid molecule with components within the test sample. Such assays will typically involve arrays comprising many genes, at least one of which is a gene of the present invention and or alleles of the protein of the present invention encoding gene of the present invention. Figure 3 provides SNP information that was identified from chromosomal sequence obtained from a number of individuals as well as SNPs found during the cDNA cloning process. These include, a6094g, t7898c, a10018c, a10791t, a12451t, t14674c, g14860c, a20133g, a1025c, t12602c, g12829t, a1442t.

Conditions for incubating a nucleic acid molecule with a test sample vary. Incubation conditions depend on the format employed in the assay, the detection methods employed, and the type and nature of the nucleic acid molecule used in the assay. One skilled in the art will recognize that any one of the commonly available hybridization, amplification or array assay formats can readily be adapted to employ the novel fragments of the Human genome disclosed herein. Examples of such assays can be found in Chard, T. *An Introduction to Radioimmunoassay and Related Techniques*, Elsevier Science Publishers, Amsterdam, The Netherlands (1986); Bullock, G. R. *et al.*, *Techniques in Immunocytochemistry*, Academic Press, Orlando, FL Vol. 1 (1982), Vol. 2 (1983), Vol. 3 (1985); Tijssen, P., *Practice and Theory of Enzyme Immunoassays: Laboratory Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers, Amsterdam, The Netherlands (1985).

The test samples of the present invention include cells, protein or membrane extracts of cells. The test sample used in the above-described method will vary based on the assay format, nature of the detection method and the tissues, cells or extracts used as the sample to be assayed. Methods for preparing nucleic acid extracts or of cells are well known in the art and can be readily be adapted in order to obtain a sample that is compatible with the system utilized.

In another embodiment of the present invention, kits are provided which contain the necessary reagents to carry out the assays of the present invention.

Specifically, the invention provides a compartmentalized kit to receive, in close confinement, one or more containers which comprises: (a) a first container comprising one of the nucleic acid molecules that can bind to a fragment of the Human genome disclosed herein; and (b) one or more other containers comprising one or more of the following: wash reagents, reagents capable of detecting presence of a bound nucleic acid.

In detail, a compartmentalized kit includes any kit in which reagents are contained in separate containers. Such containers include small glass containers, plastic containers, strips of plastic, glass or paper, or arraying material such as silica. Such containers allows one to efficiently transfer reagents from one compartment to another compartment such that the samples and reagents

are not cross-contaminated, and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers will include a container which will accept the test sample, a container which contains the nucleic acid probe, containers which contain wash reagents (such as phosphate buffered saline, Tris-buffers, etc.), and containers which contain the reagents used to detect the bound probe. One skilled in the art will readily recognize that the previously unidentified protein of the present invention encoding gene of the present invention can be routinely identified using the sequence information disclosed herein can be readily incorporated into one of the established kit formats which are well known in the art, particularly expression arrays.

#### Vectors/host cells

The invention also provides vectors containing the nucleic acid molecules described herein. The term "vector" refers to a vehicle, preferably a nucleic acid molecule, which can transport the nucleic acid molecules. When the vector is a nucleic acid molecule, the nucleic acid molecules are covalently linked to the vector nucleic acid. With this aspect of the invention, the vector includes a plasmid, single or double stranded phage, a single or double stranded RNA or DNA viral vector, or artificial chromosome, such as a BAC, PAC, YAC, OR MAC.

A vector can be maintained in the host cell as an extrachromosomal element where it replicates and produces additional copies of the nucleic acid molecules. Alternatively, the vector may integrate into the host cell genome and produce additional copies of the nucleic acid molecules when the host cell replicates.

The invention provides vectors for the maintenance (cloning vectors) or vectors for expression (expression vectors) of the nucleic acid molecules. The vectors can function in prokaryotic or eukaryotic cells or in both (shuttle vectors).

Expression vectors contain cis-acting regulatory regions that are operably linked in the vector to the nucleic acid molecules such that transcription of the nucleic acid molecules is allowed in a host cell. The nucleic acid molecules can be introduced into the host cell with a separate nucleic acid molecule capable of affecting transcription. Thus, the second nucleic acid molecule may provide a trans-acting factor interacting with the cis-regulatory control region to allow transcription of the nucleic acid molecules from the vector. Alternatively, a trans-acting factor may be supplied by the host cell. Finally, a trans-acting factor can be produced from the vector itself. It is understood, however, that in some embodiments, transcription and/or translation of the nucleic acid molecules can occur in a cell-free system.

The regulatory sequence to which the nucleic acid molecules described herein can be operably linked include promoters for directing mRNA transcription. These include, but are not limited to, the left promoter from bacteriophage  $\lambda$ , the lac, TRP, and TAC promoters from *E. coli*, the early and late promoters from SV40, the CMV immediate early promoter, the adenovirus early and late promoters, and retrovirus long-terminal repeats.

In addition to control regions that promote transcription, expression vectors may also include regions that modulate transcription, such as repressor binding sites and enhancers. Examples include the SV40 enhancer, the cytomegalovirus immediate early enhancer, polyoma enhancer, adenovirus enhancers, and retrovirus LTR enhancers.

In addition to containing sites for transcription initiation and control, expression vectors can also contain sequences necessary for transcription termination and, in the transcribed region a ribosome binding site for translation. Other regulatory control elements for expression include initiation and termination codons as well as polyadenylation signals. The person of ordinary skill in the art would be aware of the numerous regulatory sequences that are useful in expression vectors. Such regulatory sequences are described, for example, in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd. ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1989).

A variety of expression vectors can be used to express a nucleic acid molecule. Such vectors include chromosomal, episomal, and virus-derived vectors, for example vectors derived from bacterial plasmids, from bacteriophage, from yeast episomes, from yeast chromosomal elements, including yeast artificial chromosomes, from viruses such as baculoviruses, papovaviruses such as SV40, Vaccinia viruses, adenoviruses, poxviruses, pseudorabies viruses, and retroviruses. Vectors may also be derived from combinations of these sources such as those derived from plasmid and bacteriophage genetic elements, e.g. cosmids and phagemids. Appropriate cloning and expression vectors for prokaryotic and eukaryotic hosts are described in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd. ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1989).

The regulatory sequence may provide constitutive expression in one or more host cells (i.e. tissue specific) or may provide for inducible expression in one or more cell types such as by temperature, nutrient additive, or exogenous factor such as a hormone or other ligand. A variety of vectors providing for constitutive and inducible expression in prokaryotic and eukaryotic hosts are well known to those of ordinary skill in the art.

The nucleic acid molecules can be inserted into the vector nucleic acid by well-known methodology. Generally, the DNA sequence that will ultimately be expressed is joined to an expression vector by cleaving the DNA sequence and the expression vector with one or more



restriction enzymes and then ligating the fragments together. Procedures for restriction enzyme digestion and ligation are well known to those of ordinary skill in the art.

The vector containing the appropriate nucleic acid molecule can be introduced into an appropriate host cell for propagation or expression using well-known techniques. Bacterial cells include, but are not limited to, *E. coli*, *Streptomyces*, and *Salmonella typhimurium*. Eukaryotic cells include, but are not limited to, yeast, insect cells such as *Drosophila*, animal cells such as COS and CHO cells, and plant cells.

As described herein, it may be desirable to express the peptide as a fusion protein. Accordingly, the invention provides fusion vectors that allow for the production of the peptides. Fusion vectors can increase the expression of a recombinant protein, increase the solubility of the recombinant protein, and aid in the purification of the protein by acting for example as a ligand for affinity purification. A proteolytic cleavage site may be introduced at the junction of the fusion moiety so that the desired peptide can ultimately be separated from the fusion moiety. Proteolytic enzymes include, but are not limited to, factor Xa, thrombin, and enterokinase. Typical fusion expression vectors include pGEX (Smith *et al.*, *Gene* 67:31-40 (1988)), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein. Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann *et al.*, *Gene* 69:301-315 (1988)) and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185:60-89 (1990)). Recombinant protein expression can be maximized in host bacteria by providing a genetic background wherein the host cell has an impaired capacity to proteolytically cleave the recombinant protein. (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Alternatively, the sequence of the nucleic acid molecule of interest can be altered to provide preferential codon usage for a specific host cell, for example *E. coli*. (Wada *et al.*, *Nucleic Acids Res.* 20:2111-2118 (1992)).

The nucleic acid molecules can also be expressed by expression vectors that are operative in yeast. Examples of vectors for expression in yeast e.g., *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, *EMBO J.* 6:229-234 (1987)), pMFa (Kurjan *et al.*, *Cell* 30:933-943(1982)), pJRY88 (Schultz *et al.*, *Gene* 54:113-123 (1987)), and pYES2 (Invitrogen Corporation, San Diego, CA).

The nucleic acid molecules can also be expressed in insect cells using, for example, baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith *et al.*, *Mol. Cell Biol.* 3:2156-2165 (1983)) and the pVL series (Lucklow *et al.*, *Virology* 170:31-39 (1989)).

In certain embodiments of the invention, the nucleic acid molecules described herein are expressed in mammalian cells using mammalian expression vectors. Examples of mammalian expression vectors include pCDM8 (Seed, B. *Nature* 329:840(1987)) and pMT2PC (Kaufman *et al.*, *EMBO J.* 6:187-195 (1987)).

5        The expression vectors listed herein are provided by way of example only of the well-known vectors available to those of ordinary skill in the art that would be useful to express the nucleic acid molecules. The person of ordinary skill in the art would be aware of other vectors suitable for maintenance propagation or expression of the nucleic acid molecules described herein. These are found for example in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory*  
10 *Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.*

      The invention also encompasses vectors in which the nucleic acid sequences described herein are cloned into the vector in reverse orientation, but operably linked to a regulatory sequence that permits transcription of antisense RNA. Thus, an antisense transcript can be produced to all, or to a  
15        portion, of the nucleic acid molecule sequences described herein, including both coding and non-coding regions. Expression of this antisense RNA is subject to each of the parameters described above in relation to expression of the sense RNA (regulatory sequences, constitutive or inducible expression, tissue-specific expression).

      The invention also relates to recombinant host cells containing the vectors described herein.  
20        Host cells therefore include prokaryotic cells, lower eukaryotic cells such as yeast, other eukaryotic cells such as insect cells, and higher eukaryotic cells such as mammalian cells.

      The recombinant host cells are prepared by introducing the vector constructs described herein into the cells by techniques readily available to the person of ordinary skill in the art. These include, but are not limited to, calcium phosphate transfection, DEAE-dextran-mediated transfection, cationic  
25        lipid-mediated transfection, electroporation, transduction, infection, lipofection; and other techniques such as those found in Sambrook, *et al.* (*Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.*)

      Host cells can contain more than one vector. Thus, different nucleotide sequences can be introduced on different vectors of the same cell. Similarly, the nucleic acid molecules can be  
30        introduced either alone or with other nucleic acid molecules that are not related to the nucleic acid molecules such as those providing trans-acting factors for expression vectors. When more than one vector is introduced into a cell, the vectors can be introduced independently, co-introduced or joined to the nucleic acid molecule vector.

In the case of bacteriophage and viral vectors, these can be introduced into cells as packaged or encapsulated virus by standard procedures for infection and transduction. Viral vectors can be replication-competent or replication-defective. In the case in which viral replication is defective, replication will occur in host cells providing functions that complement the defects.

5        Vectors generally include selectable markers that enable the selection of the subpopulation of cells that contain the recombinant vector constructs. The marker can be contained in the same vector that contains the nucleic acid molecules described herein or may be on a separate vector. Markers include tetracycline or ampicillin-resistance genes for prokaryotic host cells and dihydrofolate reductase or neomycin resistance for eukaryotic host cells. However, any marker that provides  
10       selection for a phenotypic trait will be effective.

While the mature proteins can be produced in bacteria, yeast, mammalian cells, and other cells under the control of the appropriate regulatory sequences, cell-free transcription and translation systems can also be used to produce these proteins using RNA derived from the DNA constructs described herein.

15       Where secretion of the peptide is desired, appropriate secretion signals are incorporated into the vector. The signal sequence can be endogenous to the peptides or heterologous to these peptides.

Where the peptide is not secreted into the medium, the protein can be isolated from the host cell by standard disruption procedures, including freeze thaw, sonication, mechanical disruption, use of lysing agents and the like. The peptide can then be recovered and purified by well-known purification  
20       methods including ammonium sulfate precipitation, acid extraction, anion or cationic exchange chromatography, phosphocellulose chromatography, hydrophobic-interaction chromatography, affinity chromatography, hydroxylapatite chromatography, lectin chromatography, or high performance liquid chromatography.

It is also understood that depending upon the host cell in recombinant production of the  
25       peptides described herein, the peptides can have various glycosylation patterns, depending upon the cell, or maybe non-glycosylated as when produced in bacteria. In addition, the peptides may include an initial modified methionine in some cases as a result of a host-mediated process.

#### Uses of vectors and host cells

30       The recombinant host cells expressing the peptides described herein have a variety of uses. First, the cells are useful for producing a protein of the present invention or peptide that can be further purified to produce desired amounts of a protein of the present invention or fragments. Thus, host cells containing expression vectors are useful for peptide production.

Host cells are also useful for conducting cell-based assays involving the protein of the present invention or fragments, such as those described above as well as other formats known in the art. Thus, a recombinant host cell expressing a native protein of the present invention is useful for assaying compounds that stimulate or inhibit a protein of the present invention function.

5 Host cells are also useful for identifying mutants in which these functions are affected. If the mutants naturally occur and give rise to a pathology, host cells containing the mutations are useful to assay compounds that have a desired effect on the mutant protein of the present invention (for example, stimulating or inhibiting function) which may not be indicated by their effect on the native protein of the present invention.

10 Genetically engineered host cells can be further used to produce non-human transgenic animals. A transgenic animal is preferably a mammal, for example a rodent, such as a rat or mouse, in which one or more of the cells of the animal include a transgene. A transgene is exogenous DNA which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal in one or more cell types or tissues of the transgenic  
15 animal. These animals are useful for studying the function of a protein of the present invention and identifying and evaluating modulators of protein activity. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, and amphibians.

A transgenic animal can be produced by introducing nucleic acid into the male pronuclei of a fertilized oocyte, e.g., by microinjection, retroviral infection, and allowing the oocyte to develop in a  
20 pseudopregnant female foster animal. Any of the protein of the present invention encoding nucleotide sequences can be introduced as a transgene into the genome of a non-human animal, such as a mouse.

Any of the regulatory or other sequences useful in expression vectors can form part of the transgenic sequence. This includes intronic sequences and polyadenylation signals, if not already included. A tissue-specific regulatory sequence(s) can be operably linked to the transgene to direct  
25 expression of the protein of the present invention to particular cells.

Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Patent No. 4,873,191 by Wagner *et al.* and in Hogan, B., *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory  
30 Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the transgene in its genome and/or expression of transgenic mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene can further be bred to other transgenic animals carrying other transgenes.

A transgenic animal also includes animals in which the entire animal or tissues in the animal have been produced using the homologously recombinant host cells described herein.

In another embodiment, transgenic non-human animals can be produced which contain selected systems that allow for regulated expression of the transgene. One example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the *cre/loxP* recombinase system, see, e.g., Lakso *et al. PNAS* 89:6232-6236 (1992). Another example of a recombinase system is the FLP recombinase system of *S. cerevisiae* (O'Gorman *et al. Science* 251:1351-1355 (1991)). If a *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the *Cre* recombinase and a selected protein is required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut, I. *et al. Nature* 385:810-813 (1997) and PCT International Publication Nos. WO 97/07668 and WO 97/07669. In brief, a cell, e.g., a somatic cell, from the transgenic animal can be isolated and induced to exit the growth cycle and enter G<sub>0</sub> phase. The quiescent cell can then be fused, e.g., through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyst and then transferred to pseudopregnant female foster animal. The offspring born of this female foster animal will be a clone of the animal from which the cell, e.g., the somatic cell, is isolated.

Transgenic animals containing recombinant cells that express the peptides described herein are useful to conduct the assays described herein in an *in vivo* context. Accordingly, the various physiological factors that are present *in vivo* and that could effect substrate binding, protein activation, and signal transduction, may not be evident from *in vitro* cell-free or cell-based assays. Accordingly, it is useful to provide non-human transgenic animals to assay *in vivo* protein function, including substrate interaction, the effect of specific mutant proteins on protein function and substrate interaction, and the effect of chimeric protein of the present invention. It is also possible to assess the effect of null mutations, that is mutations that substantially or completely eliminate one or more of the proteins functions.

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred

embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the above-described modes for carrying out the invention which are obvious to those skilled in the field of molecular biology or related fields are intended to be within the scope of the following claims.

### Claims

That which is claimed is:

1. An isolated peptide consisting of an amino acid sequence selected from the group consisting of:
  - (a) an amino acid sequence shown in SEQ ID NOS:4-6;
  - (b) an amino acid sequence of an allelic variant of an amino acid sequence shown in SEQ ID NOS:4-6, wherein said allelic variant is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1-3 (transcript/cDNA) or 7 (genomic);
  - (c) an amino acid sequence of an ortholog of an amino acid sequence shown in SEQ ID NOS:4-6, wherein said ortholog is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1-3 (transcript/cDNA) or 7 (genomic); and
  - (d) a fragment of an amino acid sequence shown in SEQ ID NOS:4-6, wherein said fragment comprises at least 10 contiguous amino acids.
2. An isolated peptide comprising an amino acid sequence selected from the group consisting of:
  - (a) an amino acid sequence shown in SEQ ID NOS:4-6;
  - (b) an amino acid sequence of an allelic variant of an amino acid sequence shown in SEQ ID NOS:4-6, wherein said allelic variant is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1-3 (transcript/cDNA) or 7 (genomic);
  - (c) an amino acid sequence of an ortholog of an amino acid sequence shown in SEQ ID NOS:4-6, wherein said ortholog is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1-3 (transcript/cDNA) or 7 (genomic); and
  - (d) a fragment of an amino acid sequence shown in SEQ ID NOS:4-6, wherein said fragment comprises at least 10 contiguous amino acids.
3. An isolated antibody that selectively binds to a peptide of claim 2.

4. An isolated nucleic acid molecule consisting of a nucleotide sequence selected from the group consisting of:

(a) a nucleotide sequence that encodes an amino acid sequence shown in SEQ ID NOS:4-6;

(b) a nucleotide sequence that encodes of an allelic variant of an amino acid sequence shown in SEQ ID NOS:4-6, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1-3 (transcript/cDNA) or 7 (genomic);

(c) a nucleotide sequence that encodes an ortholog of an amino acid sequence shown in SEQ ID NOS:4-6, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1-3 (transcript/cDNA) or 7 (genomic);

(d) a nucleotide sequence that encodes a fragment of an amino acid sequence shown in SEQ ID NOS:4-6, wherein said fragment comprises at least 10 contiguous amino acids; and

(e) a nucleotide sequence that is the complement of a nucleotide sequence of (a)-(d).

5. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

(a) a nucleotide sequence that encodes an amino acid sequence shown in SEQ ID NOS:4-6;

(b) a nucleotide sequence that encodes of an allelic variant of an amino acid sequence shown in SEQ ID NOS:4-6, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1-3 (transcript/cDNA) or 7 (genomic);

(c) a nucleotide sequence that encodes an ortholog of an amino acid sequence shown in SEQ ID NOS:4-6, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1-3 (transcript/cDNA) or 7 (genomic);

(d) a nucleotide sequence that encodes a fragment of an amino acid sequence shown in SEQ ID NOS:4-6, wherein said fragment comprises at least 10 contiguous amino acids; and

(e) a nucleotide sequence that is the complement of a nucleotide sequence of (a)-(d).



6. A gene chip comprising a nucleic acid molecule of claim 5.
7. A transgenic non-human animal comprising a nucleic acid molecule of claim 5.
8. A nucleic acid vector comprising a nucleic acid molecule of claim 5.
9. A host cell containing the vector of claim 8.
10. A method for producing any of the peptides of claim 1 comprising introducing a nucleotide sequence encoding any of the amino acid sequences in (a)-(d) into a host cell, and culturing the host cell under conditions in which the peptides are expressed from the nucleotide sequence.
11. A method for producing any of the peptides of claim 2 comprising introducing a nucleotide sequence encoding any of the amino acid sequences in (a)-(d) into a host cell, and culturing the host cell under conditions in which the peptides are expressed from the nucleotide sequence.
12. A method for detecting the presence of any of the peptides of claim 2 in a sample, said method comprising contacting said sample with a detection agent that specifically allows detection of the presence of the peptide in the sample and then detecting the presence of the peptide.
13. A method for detecting the presence of a nucleic acid molecule of claim 5 in a sample, said method comprising contacting the sample with an oligonucleotide that hybridizes to said nucleic acid molecule under stringent conditions and determining whether the oligonucleotide binds to said nucleic acid molecule in the sample.
14. A method for identifying a modulator of a peptide of claim 2, said method comprising contacting said peptide with an agent and determining if said agent has modulated the function or activity of said peptide.
15. The method of claim 14, wherein said agent is administered to a host cell comprising an expression vector that expresses said peptide.

16. A method for identifying an agent that binds to any of the peptides of claim 2, said method comprising contacting the peptide with an agent and assaying the contacted mixture to determine whether a complex is formed with the agent bound to the peptide.
17. A pharmaceutical composition comprising an agent identified by the method of claim 16 and a pharmaceutically acceptable carrier therefor.
18. A method for treating a disease or condition mediated by a human drug-metabolizing enzyme protein, said method comprising administering to a patient a pharmaceutically effective amount of an agent identified by the method of claim 16.
19. A method for identifying a modulator of the expression of a peptide of claim 2, said method comprising contacting a cell expressing said peptide with an agent, and determining if said agent has modulated the expression of said peptide.
20. An isolated human proteins of the present invention having an amino acid sequence that shares at least 70% homology with an amino acid sequence shown in SEQ ID NOS:4-6.
21. A peptide according to claim 20 that shares at least 90 percent homology with an amino acid sequence shown in SEQ ID NOS:4-6.
22. An isolated nucleic acid molecule encoding a human proteins of the present invention, said nucleic acid molecule sharing at least 80 percent homology with a nucleic acid molecule shown in SEQ ID NOS:1-3 (transcript/cDNA) or 7 (genomic).
23. A nucleic acid molecule according to claim 22 that shares at least 90 percent homology with a nucleic acid molecule shown in SEQ ID NOS:1-3 (transcript/cDNA) or 7 (genomic).

## FIGURE 1, page 1 of 2

## Transcript 1:

```
1  ATTCCCAATG  GCGAAGATTG  AGAAAAACGC  TCCCACGATG  GAAAAAAGC
51  CAGAACTGTT  TAACATCATG  GAAGTAGATG  GAGTCCCTAC  GTTGATATTA
101 TCAAAAGAAT  GGTGGGAAAA  AGTATGTAAT  TTCCAAGCCA  AGCCTGATGA
151 TCTTATTCTG  GCAACTTACC  CAAAGTCAGG  TACAACATGG  ATGCATGAAA
201 TTTTAGACAT  GATTCTAAAT  GATGGTGATG  TGGAGAAATG  CAAAAGAGCC
251 CAGACTCTAG  ATAGACACGC  TTTCTTGAA  CTGAAATTTC  CCCATAAAGA
301 AAAACCAGAT  TTGGAGTTCG  TTCTTGAAAT  GTCCTCACCA  CAACTGATAA
351 AAACACATCT  CCCTTCACAT  CTGATTCCAC  CATCTATCTG  GAAAGAAAAC
401 TGCAAGATTG  TCTATGTGGC  CAGAAATCCC  AAGGATTGCC  TGGTGTCTTA
451 CTACCACTTT  CACAGGATGG  CTTCTTTTAT  GCCTGATCCT  CAGAACTTAG
501 AGGAATTTTA  TGAGAAATTC  ATGTCCGGAA  AAGTTGTTGG  CGGGTCCTGG
551 TTTGACCATG  TGAAAGGATG  GTGGGCTGCA  AAAGACATGC  ACCGGATCCT
601 CTACCTCTTC  TACGAGGATA  TTAAAAAATA  TCCAAAACAT  GAGATCCACA
651 AGGTGTTGGA  ATTCTTGAG  AAAACTTGGT  CAGGTGATGT  TATAAACAAG
701 ATTGCCACC  ATACCTCATT  TGATGTAATG  AAGGATAATC  CCATGGCCAA
751 CCATACTGCG  GTACCTGCTC  ACATATTCAA  TCACTCCATC  TCAAAATTTA
801 TGAGGAAAGG  GATGCCTGGA  GACTGGAAGA  ACCACTTTAC  TGTGGCTTTG
851 AATGAGAACT  TTGATAAGCA  TTATGAAAAG  AAGATGGCAG  GGTCCACACT
901 GAACCTCTGC  CTGGAGATCT  GAGAGGAACA  ACAACAACT  AG
(SEQ ID NO:1)
```

## FEATURES:

Start Codon: 8

Stop Codon: 920

## Transcript 2:

```
1  ATTCCCAATG  GCGAAGATTG  AGAAAAACGC  TCCCACGATG  GAAAAAAGC
51  CAGAACTGTT  TAACATCATG  GAAGTAGATG  GAGTCCCTAC  GTTGATATTA
101 TCAAAAGAAT  GGTGGGAAAA  AGTATGTAAT  TTCCAAGCCA  AGCCTGATGA
151 TCTTATTCTG  GCAACTTACC  CAAAGTCAGG  TACAACATGG  ATGCATGAAA
201 TTTTAGACAT  GATTCTAAAT  GATGGTGATG  TGGAGAAATG  CAAAAGAGCC
251 CAGACTCTAG  ATAGACACGC  TTTCTTGAA  CTGAAATTTC  CCCATAAAGA
301 AAAACCAGAT  TTGGAGTTCG  TTCTTGAAAT  GTCCTCACCA  CAACTGATAA
351 AAACACATCT  CCCTTCACAT  CTGATTCCAC  CATCTATCTG  GAAAGAAAAC
401 TGCAAGATTG  TCTATGTGGC  CAGAAATCCC  AAGGATTGCC  TGGTGTCTTA
451 CTACCACTTT  CACAGGATGG  CTTCTTTTAT  GCCTGATCCT  CAGAACTTAG
501 AGGAATTTTA  TGAGAAATTC  ATGTCCGGAA  AAGTTGTTGG  CGGGTCCTGG
551 TTTGACCATG  TGAAAGGATG  GTGGGCTGCA  AAAGACATGC  ACCGGATCCT
601 CTACCTCTTC  TACGAGGATA  TTAAAAAAGA  CCCAAAGCGG  GAAATTGAGA
651 AGATACTGAA  GTTCCTGGAA  AAAGACATAT  CAGAGGAAAT  TCTGAATAAA
701 ATCATCTATC  ACACCTCCTT  TGATGTAATG  AAGCAAAACC  CAATGACCAA
751 CTATACCACT  TTGCCACCA  GCATTATGGA  CCACTCCATC  TCCCCTTTTA
801 TGAGGAAAGG  GATGCCTGGA  GACTGGAAGA  ACTATTTTAC  TGTGGCCCAA
851 AATGAAGAAT  TTGACAAGGA  CTACCAGAAG  AAGATGGCAG  GAAGCACCTT
901 AACCTTCCGC  ACAGAGATCT  GA
(SEQ ID NO:2)
```

## FEATURES:

Start: 8

Stop: 920

2/16

## FIGURE 1, page 2 of 2

## cDNA sequence

```

1   CTCACTATTA GGGCGAATTG AATTTAGCGG CCGCGAATTC GCCCTTATGG
51  CGAAGATTGA GAAAAACGCT CCCACGATGG AAAAAAGCC AGAACTGTTT
101 AACATCATGG AAGTAGATGG AGTCCCTACG TTGATATTAT CAAAAGAATG
151 GTGGGAAAAA GTCTGTAATT TCCAAGCCAA GCCTGATGAT CTTATTCTGG
201 CAACTTACCC AAAGTCAGGT ACAACATGGA TGCATGAAAT TTTAGACATG
251 ATTCTAAATG ATGGTGATGT GGAGAAATGC AAAAGAGCCC AGACTCTAGA
301 TAGACACGCT TTCCTTGAAC TGAAATTTC CCATAAAGAA AAACCAGATT
351 TGGAGTTCGT TCTTGAAATG TCCTCACCAC AACTGATAAA AACACATCTC
401 CCTTCACATC TGATTCCACC ATCTATCTGG AAAGAAAAC GCAAGATTGT
451 CTATGTGGCC AGAAATCCCA AGGATTGCCT GGTGTCCTAC TACCACTTTC
501 ACAGGATGGC TTCCTTTATG CCTGATCCTC AGAACTTAGA GGAATTTTAT
551 GAGAAATTCA TGTCCGGAAA AGTTGTTGGC GGGTCCTGGT TTGACCATGT
601 GAAAGGATGG TGGGCTGCAA AAGACACGCA CCGGATCCTC TACCTCTTCT
651 ACGAGGATAT TAAAAAAAT CCAAACATG AGATCCACAA GGTGTTGGAA
701 TTCTTGGAGA AAACCTTTGTC AGGTGATGTT ATAAACAAGA TTGTCCACCA
751 TACCTCATTT GATGTAATGA AGGATAATCC CATGGCCAAC CATACTGCGG
801 TACCTGCTCA CATATTCAAT CACTCCATCT CAAAATTTAT GAGGAAAGGG
851 ATGCCTGGAG ACTGGAAGAA CCACTTTACT GTGGCTATGA ATGAGAACTT
901 TGATAAGCAT TATGAAAAGA AGATGGCAGG GTCCCACTG AACTTCTGCC
951 TGGAGATCTG AGAGGAACAA CAAAGGCGA ATTCGTTTAA ACCTGCAGGA
1001 CTAG

```

(SEQ ID NO:3)

## FEATURES:

5'UTR: 1-47  
 Start: 47  
 Stop: 959  
 3'UTR: 959-1004

## Homologous proteins:

## Top 10 BLAST Hits

	Score	E
gi 1711569 sp P50237 SUAC_RAT N-HYDROXYARYLAMINE SULFOTRANSFERA...	388	e-107
gi 3004922 gb AAC17740.1  (AF033653) phenol sulfotransferase [M...	388	e-107
gi 8117877 gb AAF72810.1 AF186263_1 (AF186263) sulfotransferase...	360	2e-98
gi 5730071 ref NP_006579.1  SULT1C sulfotransferase >gi 3649608...	360	2e-98
gi 4507305 ref NP_001047.1  sulfotransferase family 1C, member ...	348	4e-95
gi 4689040 emb CAB41460.1  (AJ238391) sulfotransferase K1 [Ratt...	344	6e-94
gi 2828826 gb AAC00410.1  (AF026304) sulfotransferase [Oryctola...	340	1e-92
gi 4689042 emb CAB41461.1  (AJ238392) sulfotransferase K2 [Ratt...	338	6e-92
gi 2290540 gb AAB65154.1  (U95726) thyroid hormone sulfotransfe...	332	3e-90
gi 4096440 gb AAC99889.1  (U32371) tyrosine-ester sulfotransfer...	332	4e-90
gi 7657621 ref NP_055280.1  thyroid hormone slfotransferase >gi...	332	4e-90
gi 7949146 ref NP_058051.1  amine N-sulfotransferase >gi 743496...	329	3e-89
gi 8117857 gb AAF72802.1 AF186254_1 (AF186254) sulfotransferase...	328	6e-89

## EXPRESSION INFORMATION FOR MODULATORY USE:

library source for cDNA:  
 Human Kidney  
 Human Small Intestine

3/16

## FIGURE 2, page 1 of 5

Protein 1 (from Transcript 1):

```

1  MAKIEKNAPT MEKKPELFNI MEVDGVPTLI LSKEWWEKVC NFQAKPDDLI
51 LATYPKSGTT WMHEILDMIL NDGDVEKCKR AQTLDRAHFL ELKFPKKEKP
101 DLEFVLEMSS PQLIKTHLPS HLIPPSIWKE NCKIVYVARN PKDCLVSYH
151 FHRMASFMPD PQNLEEFYEK FMSGKVVGGS WFDHVKGWWA AKDMHRILYL
201 FYEDIKKNPK HEIHKVLEFL EKTWSGDVIN KIVHHTSFDV MKDNPMAHNT
251 AVPAHIFNHS ISKFMKRGMP GDWKNHFTVA LNFNFDKHYE KKMAGSTLNF
301 CLEI

```

(SEQ ID NO:4)

**Features****Hummer Results:**

Scores for sequence family classification (score includes all domains):

Model	Description	Score	E-value	N
PF00685	Sulfotransferase proteins	512.4	3.2e-150	1

Parsed for domains:

Model	Domain	seq-f	seq-t	hmm-f	hmm-t	score	E-value
PF00685	1/1	24	292 ..	1	281 []	512.4	3.2e-150

**Important Domains****Prosite search results:**

[1] PDOC00001 PS00001 ASN\_GLYCOSYLATION

N-glycosylation site

Number of matches: 2

1	248-251	NHTA
2	258-261	NHSI

[2] PDOC00005 PS00005 PKC\_PHOSPHO\_SITE

Protein kinase C phosphorylation site

173-175 SGK

[3] PDOC00006 PS00006 CK2\_PHOSPHO\_SITE

Casein kinase II phosphorylation site

Number of matches: 2

1	180-183	SWFD
2	236-239	TSFD

[4] PDOC00008 PS00008 MYRISTYL

N-myristoylation site

295-300 GSTLNF

## FIGURE 2, page 2 of 5

Alignment to Top Blast Hit:

>gi|1711569|sp|P50237|SUAC\_RAT N-HYDROXYARYLAMINE SULFOTRANSFERASE  
(HAST-I) >gi|543420|pir||A49098 N-hydroxyarylamine  
sulfotransferase, HAST-I - rat >gi|440841|gb|AAA42181.1|  
(L22339) N-hydroxy-2-acetylaminofluorene [Rattus  
norvegicus]  
Length = 304

Score = 388 bits (987), Expect = e-107

Identities = 174/283 (61%), Positives = 220/283 (77%)

Frame = +2

Query: 71 EVDGVPTLI LSKWEWEKVCNFQAKPDDLILATYPKSGTTWMHEILDMILNDGDVEKCKRA 250  
EV+G+ L + W+K+ NFQAKPDDL++ATY K+GTTW EI+DMI NDGDV+KC+RA  
Sbjct: 22 EVNGILMSKILMSDNWDKIWNFQAKPDDLIIATYAKAGTTWTQEIIVDMIQNDGDVQKQORA 81  
Query: 251 QTLDRHAFLELKFPHEKPDLEFVLEMSSPQLIKTHLPSHLIPPSIWKENCKIVYVARNP 430  
T DRH F+E P L+ +M SP+ +KTHLP H++PPS WKEN KI+YVARN  
Sbjct: 82 NTYDRHPFIEWTLPSPLNSGLDLANKMPSPTLKTLPVHMLPPSFWKENSKIYYVARNA 141  
Query: 431 KDCLVSYHFFHRMASFMPDPQNLEEFYEKFMGKVVGGSWFDHVKGWAAKDMHRILYLF 610  
KDCLVSYH+F RM +PDP L E+ E+F +GKV+ GSW+DHVKGW KD HRILYLF  
Sbjct: 142 KDCLVSYHFFSRMNKMLPDPGTLGEYIEQFKAGKVLWGSWYDHVKGWWDVQDHRILYLF 201  
Query: 611 YEDIKKNPKHEIHKVLEFLEKTWSGDVINKIVHHTSFDVMKDNPMANHTAVPAHIFNHSI 790  
YED+K++PK EI K+ +FLEK S +V+NKI++HTSFDVMK+NPMAN+T +P+ I +HSI  
Sbjct: 202 YEDMKEDPKREIKKIAKFEKDISSEVLNKIIYHTSFDVMKENPMANYTTLPSSIMDHSI 261  
Query: 791 SKFMRKGMPPGDWKNHFTVALNENFDKHYEKKMAGSTLNFCEI 919  
S FMRKGMPPGDWKN+FTVA +E+FD+ Y +KMAGS + F EI  
Sbjct: 262 SPFMRKGMPPGDWKNYFTVAQSEDFEDYRRKMAGSNITFRTEI 304  
(SEQ ID NO: 8)

Membrane Spanning Regions:

None found

Protein 2 (from Transcript 2):

1 MAKIEKNAPT MEKKPELFNI MEVDGVPTLI LSKWEWEKVC NFQAKPDDLI  
51 LATYPKSGTT WMHEILDMIL NDGDVEKCKR AQTLDRAHFL ELKFPHEKCP  
101 DLEFVLEMSS PQLIKTHLPS HLIPPSIWKE NCKIVYVARN PKDCLVSYH  
151 FHRMASFMPD PQNLEEFYEK FMSGKVVGGG WFDHVKGWAA AKDMHRILYL  
201 FYEDIKKDPK REIEKILKFL EKDISSEILN KIIYHTSFDV MKQNPMTNYT  
251 TLPTSIMDHS ISPFMRKGMPP GDWKNYFTVA QNEEFDKDYQ KKMAGSTLTF  
301 RTEI  
(SEQ ID NO:5)

Features:Hmmer Results:

Scores for sequence family classification (score includes all domains):

Model	Description	Score	E-value	N
PF00685	Sulfotransferase proteins	550.5	1.2e-161	1

Parsed for domains:

Model	Domain	seq-f	seq-t	hmm-f	hmm-t	score	E-value
PF00685	1/1	24	292	1	281	550.5	1.2e-161

Membrane Spanning Regions:

None found

## FIGURE 2, page 3 of 5

Important Domains:Prosite search results:

[1] PDOC00001 PS00001 ASN\_GLYCOSYLATION  
N-glycosylation site  
248-251 NYTT

---

[2] PDOC00005 PS00005 PKC\_PHOSPHO\_SITE  
Protein kinase C phosphorylation site  
Number of matches: 2  
1 173-175 SGK  
2 299-301 TFR

---

[3] PDOC00006 PS00006 CK2\_PHOSPHO\_SITE  
Casein kinase II phosphorylation site  
Number of matches: 3  
1 180-183 SWFD  
2 236-239 TSFD  
3 255-258 SIMD

---

[4] PDOC00008 PS00008 MYRISTYL  
N-myristoylation site  
295-300 GSTLTF

Alignment to Top Blast Hit:

>gi|1711569|sp|P50237|SUAC\_RAT N-HYDROXYARYLAMINE SULFOTRANSFERASE  
(HAST-I) >gi|543420|pir||A49098 N-hydroxyarylamine  
sulfotransferase, HAST-I - rat >gi|440841|gb|AAA42181.1|  
(L22339) N-hydroxy-2-acetylaminofluorene [Rattus  
norvegicus]  
Length = 304

Score = 435 bits (1107), Expect = e-121  
Identities = 196/283 (69%), Positives = 234/283 (82%)  
Frame = +2

Query: 71 EVDGVPTLILSKEWKEVCNFOAKPDDLILATYPKSGTTWMHEILDMILNDGDVEKCKRA 250  
EV+G+ L + W+K+ NFQAKPDDL++ATY K+GTTW EI+DMI NDGDV+KC+RA  
Sbjct: 22 EVNGILMSKILMSDNWDKIWNFOAKPDDLILATYAKAGTTWTQEIIVDMIQNDGDVQKCQRA 81  
Query: 251 QTLDRHAFLELKFPKHEKPDLEFVLEMSSPOLIKTHLP SHLIPPSIWKENCKIYVARNP 430  
T DRH F+E P L+ +M SP+ +KTHLP H++PPS WKEN KI+YVARN  
Sbjct: 82 NTYDRHPFIEWTLPSPLNSGLDLANKMPSRPTLKT HLPVHMLPPSFWKENS KIIYVARNA 141  
Query: 431 KDCLVSYHFFHRMASFMPDQNLLEEFYEKFMGKVVGGSWFDHVKGWAAKDMHRILYLF 610  
KDCLVSYH+F RM +PDP L E+ E+F +GKV+ GSW+DHVKGWW KD HRILYLF  
Sbjct: 142 KDCLVSYHFFSRMNKMLPDPGTLGEYIEQFKAGKVLGWSWYDHVKGWWDVQDHRILYLF 201  
Query: 611 YEDIKKDPKREIEKILKFLEKDISEEILNKIIYHTSFDVMKQNPMTNYTTLPTSIMDHSI 790  
YED+K+DPKREI+KI KFLEKDISEE+LNKIIYHTSFDVMK+NPM NYTTLP+SIMDHSI  
Sbjct: 202 YEDMKEDPKREIKKIAKFLEKDISEEVLNKIIYHTSFDVMKENPMANYTTLPSIMDHSI 261  
Query: 791 SPFMRKGM PGDWKNYFTVAQNEEFDKDYQKKMAGSTLTFRTEI 919  
SPFMRKGM PGDWKNYFTVAQ+E+FD+DY++KMAGS +TFRTEI  
Sbjct: 262 SPFMRKGM PGDWKNYFTVAQSEDFEDYRRKMAGSNITFRTEI 304  
(SEQ ID NO: 9)

6/16

## FIGURE 2, page 4 of 5

Protein 3 (from cDNA):

```

1  MAKIEKNAPT MEKKPELFNI MEVDGVPTLI LSKEWWEKVC NFQAKPDDLI
51  LATYPKSGTT WMHEILDMIL NDGDVEKCKR AQTLDRAHFL ELKFPKKEKP
101 DLEFVLEMSS PQLIKTHLPS HLIPPSIWKE NCKIVYVARN PKDCLVSYH
151 FHRMASFMPD PQNLEEFYEK FMSGKVVGGS WFDHVKGWWA AKDTHRILYL
201 FYEDIKKNPK HEIHKVLEFL EKTLSGDVIN KIVHHTSFDV MKDNPMANHT
251 AVPAHIFNHS ISKFMKGMP GDWKNHFTVA MNENFDKHYE KKMAGSTLNF
301 CLEI
(SEQ ID NO:6)

```

Features:Alignment to Top Blast Hit:

```

>gi|3004922|gb|AAC17740.1| (AF033653) phenol sulfotransferase [Mus
musculus]
Length = 304

```

```

Score = 391 bits (993), Expect = e-108
Identities = 175/283 (61%), Positives = 220/283 (76%)
Query: 22  EVDGVPTLILSKWEKVCNFQAKPDDLILATYPKSGTTWMHEILDMILNDGDVEKCKRA 81
          EV+G+   + E W+K+ NFQAKPDDL++ATY K+GTTW EI+DMI NDGDV+KC+RA
Sbjct: 22  EVNGILMSKMMSENWDKIWNFQAKPDDLLIATYAKAGTTWTQEIIVDMIQNDGDVQKCQRA 81
Query: 82  QTLDRHAFLELKFPKKEKPDLEFVLEMSSPQLIKTHLPSHLIPPSIWKENCKIVYVARNP 141
          T DRH F+E   P      L+   +M SP+ +KTHLP ++PPS WKEN +I+YVARN
Sbjct: 82  NTYDRHPFIEWTLPPPLNSGLDLANKMPSPRTLKTHLPVQMLPPSFWKENSQIIYVARNA 141
Query: 142 KDCLVSYHHFHRMASFMPDPQNLEEFYEKFMMSGKVVGGSWFDHVKGWWAAKDTHRILYLF 201
          KDCLVSYH+ F RM   +PDP L E+ E F +GKV+ GSW+DHVKGWW KD HRILYLF
Sbjct: 142 KDCLVSYHHFHRMASFMPDPQNLEEFYEKFMMSGKVVGGSWFDHVKGWWAAKDTHRILYLF 201
Query: 202 YEDIKKNPKHEIHKVLEFLEKTLSGDVINKIVHHTSFDVMKDNPMANHTAVPAHIFNHSI 261
          YED+K++PK EI K+++FLEK +S +V+NKI+HHTSFDVMK NPMAN+T +P+ I +HSI
Sbjct: 202 YEDMKEDPKREIKKIVKLEKDISEVLNKKIHHHTSFDVMKQNPANYTTLPSSIMDHSI 261
Query: 262 SKFMRKGMPGDWKNHFTVAMNENFDKHYEKKMAGSTLNFNCLEI 304
          S FMRKGMPGDWKN+FTVA +E+FD+ Y KKMAGST+ F EI
Sbjct: 262 SPFMRKGMPGDWKNYFTVAQSEDFEDYRKKMAGSTITFRTEI 304
(SEQ ID NO: 10)

```

Membrane Spanning Regions:

None found

Important Domains:Prosite:

[1] PDOC00001 PS00001 ASN\_GLYCOSYLATIONN-glycosylation site

Number of matches: 2

```

1  248-251 NHTA
2  258-261 NHSI

```

-----[2]  
PDOC00005 PS00005 PKC\_PHOSPHO\_SITEProtein kinase C phosphorylation site

Number of matches: 2

```

1  173-175 SGK
2  194-196 THR

```

-----[3]  
PDOC00006 PS00006 CK2\_PHOSPHO\_SITECasein kinase II phosphorylation site

Number of matches: 2

```

1  180-183 SWFD
2  236-239 TSFD

```

-----[4]  
PDOC00008



7/16

## FIGURE 2, page 5 of 5

PS00008 MYRISTYLN-myristoylation site  
295-300 GSTLNF

Hmmer:

Scores for sequence family classification (score includes all domains):

Model	Description	Score	E-value	N
PF00685	Sulfotransferase proteins	518.6	4.7e-152	1

Parsed for domains:

Model	Domain	seq-f	seq-t	hmm-f	hmm-t	score	E-value
PF00685	1/1	24	292 ..	1	281 []	518.6	4.7e-152

8/16

## FIGURE 3, page 1 of 9

```

1  AGTTCCAAAC TTTCCACAT TTTCTGTCT TCTTCTGTTT CAATCTCTGC
51  CTGTTACCCA GTTCCAAAGT CAATTTTACA TTTTCAGGTA TCTGCAGTAG
101 CACCCCACTC TGCTAGTACA AATTTATTGT ATTAGTCCCT TTTTCATGCTG
151 TTAATAAAGA CATACCTGAG ACTGGGCAAT TTAAAAAATA ACGAGGTTTA
201 ACAGACTTAA AGTTCCACGT ATCTGGGGAA GCCTCACAGT CATGGTGGAA
251 GGCAATGAAT GGCAGCAGGC AAAGGAAAAG AGAGCTTGTG CAGGGGAACT
301 CCCCTTATA ATGGGTGGGG ACACAGGCGA ACCATATCAG ACCCCAACTA
351 AATTCCAAGT TTCCAGAGTT AGTAGACATG AGAGTTTTCA GGTACATGGG
401 TACAAGAGAG TTTTTTCTGC TGTAACCCAT GACTGATATT TCAAAAATTA
451 TTCCATGAAT GAAAAAATA CTACATGAAA TTATGTTTTT TGAAGATTTT
501 CTCTGTGAAA ACTATTCAAG AAAATTGAGT ATAGAATGCT CCTTAAACC
551 ATTGTTTTGA ATTTTCTTCA ATGTAATTGT CTCGTTCTA ATTATACAAA
601 ATAATATCTT GAAGACAATG AAGCAATATA TGACACATAA CCCATTATAT
651 TCCCGATTAA CAAGTAATAT GGTGTAGG TCATGCAGCA GGAAAAAGAC
701 TATTGGGTA TAAATCTCAG CCCCAACAGG TATTGATTAT TTTACTTTTA
751 TCAAATTATT TAAACTTATA ATGACTTAAG TTTCCACATG TAGAAAACAG
801 AAAAAACTAT TACCTATTTT ATGTGTTGTC TATGAAGATT AAATAATTAA
851 TGTACATAGA GTAAGTTGTT AGCATGTTAT ATGTTAGCTT TCACTTATTT
901 TATGCTATTC TAATCAAAGG CAGTAAGATG ATAACCCATA CTATGAACCA
951 TAAAACAACA TTTTAAAAA TTTTAATTAG TATTGATCTT ACCCATCCCA
1001 GATTCCCAAT GGCGAAGATT GAGAAAAACG CTCCCACGAT GGAAAAAAG
1051 CCAGAACTGT TTAACATCAT GGAAGTAGAT GGAGTCCCTA CGTTGATATT
1101 ATCAAAAGAA TGGTGGGAAA AAGTATGTAA TTTCCAAGCC AAGCCTGATG
1151 ATCTTATTCT GGCAACTTAC CCAAAGTCAG GTAAGGGTAG CAAAACATAA
1201 AAATATTCAA TATTTTCACG TGAATTATT GCATAATCTG TATTGATAAA
1251 TGAAGCATGA TTGGGATTG GAGAGAAACA ATTCCTCATI ATGGAGATCT
1301 GTTCTTTGGT GCTGCAGGAC ATTTAGCATT CCTGATCCCT AGGACAACAA
1351 TTTCCAATAG CACTCTCTGA AGAGAACAGA AAAGATTTCT AAATGCTTTG
1401 GTAGTAGGGC AGATAATGCT CCCATTCCAG AACCATAGC ATACATCAAA
1451 TATAAGGTCA ACAAATTGAA AGACCTATGG CTTTTTAAA CATCAACCT
1501 CAAGACAGCC TTCTAAAAGC ATGTCCTACT CCAAATATTA ATCTATTATC
1551 TCAGATATTA AACACAAATT GATTTTCTAA TCCTCTAAAG CTCTTGGAGG
1601 ATGTCAGCCA TGAATTTTCT GCTCTGTACC AAATTAGTCT CATTGAGAAA
1651 AGCCAATGA CTGACCTGA TTTAAAATGC CTTAGATTTC TAATCTTCT
1701 CTAAAAATTC CTAGACTGGA ACACATGCTA GAGTCAATGG GCACACCTAA
1751 TGCCAGAAC TCACTTTCGA TAGGCCATTG TCCATTAAAA TGAACCACAG
1801 CTATTAGGAG AATTGATGAT CCCACTCTTG GGAATAGAAT ATGCAAGGGG
1851 AATGTACTAC ATCTTCTTTT TGAAGGAGTA AGTGATCAGT ATATATTCCC
1901 AGAATTGTTT GTTTGTATTA AATGTCTGTA AAAATAAAGC ACACATAATA
1951 ACAAAAAAAT TGATGGGGAC ATGTAGAAGG ATACAAGAAC CAACGTGAAG
2001 GGGCTCTCAC AGTAGCCACA TTTGGGACAG TTTGAGCATC CAAAAGAATG
2051 ATCACTAAT GATTGGGAGA ACATTAAATA AAAACCTCCT GGTCAGCCAT
2101 GCCGAAGGGA CAAGATTGGA TGTGTACTCC AGCCATAATA AAAAAGGAAA
2151 CTGGATAAAA TATATGAAAC CACTGGTTTC TGACCTTGGA TGACAGTTGT
2201 CTCAATCCTC AGACTGAAGA TGCTTTCTGA ATAGGGTATA GTTATAGGGT
2251 GCAAAAGAAA AAGCCATAGC AAACCTATTG AGTTCAAGAG CCAGGTGTAG
2301 GAGTTCGAG AGGTTGAAGT GGCTGCTATT TGCAGGTAAG AGTACCAGAC
2351 AGAAGGAAAC TACACAAATA AATACCTTTA GATATCTTTA GAGAGGACCC
2401 TTTTAGTTTA TTGTTGAATG GTAGACTGCA CTTCCATAGA GCCTATAGCT
2451 CCATGAGATC AGGCAAGGAA CCACCAGAAA ACTATTAGCC ATATAATTCC
2501 TAGGATTAT ACAAGCATGA GAGACTTTTG TGCTCCAATC AGGATAGGAT
2551 GGACAGAATT TTGGTCCCA TGACATTAGT CCTCTGTTAT TACATCTGCC
2601 GTTATTTTCA ATTACATTCC CAAAAGGATT TTGCAGTTGC TATTAACTA
2651 TCTAATCAGC TGATATTAAA ATAGGGAGAT TATTCTGGAC TATCTTGACG
2701 GACCCAGTGG AATCATGTGA ACAGGAAGTC AGAGAGATGT GGCAGAGGAG
2751 AAAGTCTGAG AAATTTAAAG TATAATAAAA GTTCACTGCA TGATTATAGA
2801 TTTAATGATG AAGAGAGAAA GTATCAACAA AACAGTTATC TCAATACTAC

```

9/16

## FIGURE 3, page 2 of 9

```

2851 AGCCACAAGC AACTGAATTC TGTGGGCATC TAGGAGCTTG GCAAAACACC
2901 CTGAAGTCCC AGATGAGAAT GGCAGCCCTA GCTGATACCT TGATTTTAGC
2951 CTAGTGAGAC CCTAAACAGA GGACTAGCCA TGTTAACCCC AATTTCTAAT
3001 CTACAGAAAC TATGACCGAA TATTCAGGTG TTGTTTTAAC TCATGAAGCT
3051 TGTGGTAATT TGTTAACCAC AAAGTCTTCA ACCTAGAATT CCATATACCC
3101 AGTAAAAATA ATATATTAAA ATATTACTAA AAATAAAGGT CAAAGACTTT
3151 TTCTCTGACA GCGAAAAGCT GAATGTGTTG CCCGCACAGC TGCCTAAAAA
3201 ATAAATAAAT AAAAAGTTAT ATTAAAGAAA TTTATTCAGA GTAACAGAAA
3251 ATAATAGTAT AAAAACTTT ATTATCTTTC AACATTCCAG CAACACACAG
3301 TTGGAATATG ACATTTTAAA AATACCATTG ATAAAAATAA TAGCACCTGA
3351 AAACATGAAA TATTCAGAAA TGAACCTAAC AAAGCTGTGC ATGATTTGAA
3401 AGTGAAAACA CTACGGAGAG AAATTGTATA AGACTTAAAT AAATAAGGAG
3451 GCATACCCCTA TCTACGGATC AAAAGGATCA ATATTTTAA GATATCAACT
3501 GTTTCAAAAT TAATGTATAA ATTCAATCTC AATCACAAC TGACAAATTG
3551 ATTCCAAGAT TTCTATGAAA ATGTCATTAA AAGCAAATAT TTTTGATAGA
3601 GTAAAGTGGC AGGGTTTATA TTAAGTATT TCATGACATT TAACTCAGCT
3651 TCAACATTCA AAAGACTGTG ATACTGTTGA TATTGTTGAT AGACATAAAT
3701 ACTTGACCCT TACTTGGTGC CAGATGCAAA AAAGTCAAAG TGCAATGCAT
3751 CAGACCTCAA AAAATTAAAA TTAAATCTCT AAAACTATGG AAGTATAGAT
3801 AGGAGAATAG CTTCAAACCT TAGGTTAGGC AACAATAATT TGGGGAAAAA
3851 AATGAAAAGC ACTACCCTTT ATGGTTTGCA TTTAATGTAA ATTCAATATA
3901 AATTAGACTT AATCAAATTA AAAGTCTGT TCTTCTAAAG ACCCAGTTAA
3951 GAAAAATAAA ATATGTGACA TAGATGGAGA GAAAATATTC ACAATACATA
4001 TATCTGGCCA GAAGGTATAA AGAACTGTTA CAACTAAGAA CAAAAACAA
4051 AAAAAAATG TATTAAAGTG GGCAAAAGAT GTAAAAATGT ATCACCACAA
4101 AAGCTTTACT ATTAACCAAT AAACACAAGA TATTCAACAT CATTTATCAT
4151 GAGAAAATGT AAATTAGTAC TATAATGAAA TACCACTACA CACCACTTCA
4201 AATGGCTAAA AACCTGAAA TACTAAGTGT TGATAAGGAC ACAGAGCAAC
4251 TGAAATTCTC ATGAAGTTT GGGGAGGACC TAACATGGTA CTAATGTTGA
4301 ACAGTTTGT ATAAACTAA ACATGCACCT ACCACACAAC CCAGCAACTC
4351 CAATCCTAGA GATTCACCCC CACCCCAAA ATAGAAATCT ATGTTTCATAT
4401 AAAAAGTGT ACACAACACT ATAGCAGACT TATTCAAAAT TAGCCCAACA
4451 CTGAAGAAAA CCTAAATTCG TATTGTCATA TAAATGAATA AACAAATCAC
4501 TGCTTATCCA TACAATTGAA CTGTTCTCAG CAATTAAGAA AAATGAACCTA
4551 CAGATATATA CAAGAACATA AATGAATCCC AAAATAATGA GGGTGTGTGA
4601 AAAAATTCAG ACACATGAGG ACAAAGTGA TGGTTTCATT TATATGAAAT
4651 TCAACAATGT GTAAACTAA TTTGTATTGA CAAAAACATA GATCAGCAGT
4701 TCCGTGGAGT CGGAAGTGAG GAAAAGAATT ACTGATAGCA ACAGACACAA
4751 CGTTCTACTC TTATCTATGG TAATGGCTAC ATACATTTGT CAAATTTCTT
4801 TAACTACAC ATTTATAACA GGTTCATTCT ATTCTATGTA ATTTTACCA
4851 CACTGAAGTT TATTTCAAAA AACGTGATTC CATAATGATG AAAAATACAA
4901 AAAGAAAAAC TTATGTATTA TGATTGAAGA TAAGTGTAT TCATCTCTTA
4951 GACTAAAAAG AAGTAATTAA GAGAAAGAAT TTAGAGGAAT TGCAGTTCTT
5001 CCCTGATTTA TGAGGGAAG TTCTTTATAT GAAGATCTAC CTAATAAATA
5051 GAGAAGTGAG GGGATTAGAA AATAAGCAAT TTGAAACCCA CAATAAAAAAT
5101 TAGGTAAAGT AGGATCATT AATGAATGA CTTAAATAA TTAGATAAAA
5151 TATGAGGAGA ACTGGCTGGT CACATGGTAC TGAAATGTCA CCACATAGTT
5201 TTATTCCTAA AAGCAAATGG AATATGGGC CTTTACAATG AGCGACCTGG
5251 TGGTCACCTA CAAAAGTGA TGATGAACAG TAGTATCACT AGTAGTGAGG
5301 CAATCAGACG TATGTACTTC TTGGCATGGT GCAAGAAGTA CCAAGCACTG
5351 CCTGAGAAAT CTGCCAAAAA TGTTCGCTAG AACCTAATCA AGACCTGGAT
5401 TCCAGTTTAT GGGAAATAGA GGAAATAGAG GAACAAAAAT GTACTGTGAA
5451 GAAATAATCA GACAAATCCA GAATGTGGG AGTACTATAG GACAGCTGGC
5501 TAATTCGTTT AAAAACC AATGTCAAAGA AAAAAATCAGT ATCATTAAAG
5551 AAAATATATG AAAGATGATT CTAAGCTGAG AAAAAAGGA TAGCAAAAGT
5601 AATGCAGCCA ATACATCTTA ATTGGGCAAA AGTGATTAAT AACATTATGA
5651 TACAATGGAA AAAAATTAAC ATAGGTTGAG TATCTCATGT TATTAAGGAT

```

10/16

## FIGURE 3, page 3 of 9

```

5701 TTATTATAAA CTAGATGTGA TGAGAAACAT TTAGTAACCA TGATAAGAGA
5751 ATGTCATATT GGAAGATACA GCATTTTGG CATGAAGTGT CATAATATAG
5801 GAAATGTAGA AATGGTTCAC TCAAACATGA TAGTTGAGAG ATTAAATACT
5851 AAAAGATGAT AGATAGATAG ACAGATAATA GGAAGATAGT AGATACATAC
5901 ATACATACAT ACTTAGATAC ATAGATAGAT ACATAGATAC ATAAAGAAAC
5951 AGATGATAGA TTAGATAGAT AGATAGATAG ATAGATAGAT AGATAGATAG
6001 ATGGATAGAT AGATGGATAG ATAGATGCAG GTATAATACA ACAGTTTTCA
6051 TTTCTGCCTT GGCAACATTG ATCTAATTTT AAAGTGCTCT CATGTTGCTG
6101 TCTTCTTCA AAATATTCAA TAACTTAGAA TGAAGTTCA ATTGACTAAA
6151 ATTAAGAAGC TATTTCAAAT ATTTTCAGGT ACAACATGGA TGCATGAAAT
6201 TTTAGACATG ATTCTAAATG ATGGTGATGT GGAGAAATGC AAAAGAGCCC
6251 AGACTCTAGA TAGACACGCT TTCCTTGAAC TGAATTTTCC CCATAAGAA
6301 AAACCAAGTG AGTAATATGC ACGAAGATAG AAAGGACTTT CACTTCAGGA
6351 TTCCAGAGCA ATGTGTACTG TCTCTGATAG AGCATCCGTG GTAGCCAGAA
6401 GTAGCCTGTA TCTTTCATGA GGTCTATTTG TTCTCAGGCC AACAATCAAA
6451 CTCTAGATTG GGTCTTCAGG GCTTCTCTCT GTTCTTAGCT GGTGGCCTTT
6501 ATCTCCCCAA TAAGATTTTC ATTCTCTTTC TCATATTTCT CCTACCCAAT
6551 GTTACGATAA AGAAGACTCC CTCTGCTTTG TATTCTCCTT CATATGTTG
6601 AATCAGTGGA AAGGCCAGAA ATTAGCAATC AGCTATTGTA AAATAACAGG
6651 TGTGATTTCT GAAGAAAAG GAGGAGAAGT AGGGGGGATA TACCGTGGAC
6701 ATTCCAGAA TTAATTGGGG GAGCTGAAA GGTTCCTCA GAGTGTA AAA
6751 CCCAGTGAAT AACATGAAAT AAAGACAGAC CTCCTAGAGG TCTTCTTTC
6801 CAAGAATGAC AACCTATTGT AAAAAATGG ATGTAATTTT TGACAAACAA
6851 AAGAGAGAGA GAAAAAGAGA AAGGAAAAA ACAACCAGAT ATATAACAAA
6901 CATTCCGATA ATTAATCGAG GGTGAAAAAG AGCTTTCCTG AAAGTCTAAA
6951 AGCTGAGAGA ATAAGAGGAA ATAAGCAAAA CTCTCCTAGA AGGTTAGGCA
7001 GGATTGGATG CTTTGTATGT CCTTGTGGAT AAAAAATTAC TATTTCCAAT
7051 GGATATAAAA CAAAATATAA TCACTCTACA GACAAAGATA TAAATGGAAT
7101 TCAAGTATTT TGGCAGAGTG CCAAGATAA ACAAAGAATA TAAATTGTTT
7151 CTTGCTAGAT TTGGAGTTCG TTCTTGAAAT GTCCTCACCA CAACTGATAA
7201 AAACACATCT CCCTTCACAT CTGATTCCAC CATCTATCTG GAAAGAAAAC
7251 TGCAAGGTAT AAAGAGGGGG CTTTTCAAAC TTCTCTTAGC TTGGTGATAT
7301 AAACATACA ACTGAAGATA TCTTTCAAAA TAATATACTT TGAAAAATAT
7351 TTTCCAAAAT ATAATTTGCT ATTTTCTTA GATGAAGCAC TTAATCAA
7401 GGATTACATA AATTTGAAAT CTGCAAACAT CCATGTTTTT TAAATTCAT
7451 TTTCTCTTAA TCCTATTTCA TGAAAAATC TTGGTAAGAT TTTCCAAAAT
7501 TGAGTCTGTG TTGCTACAAA TCAGAGAGTG TATTGGGAGC TAGAATGGAG
7551 GGAACACAT TTAATAATAA AACCTGTTG CTTGCTCTG CCAAAGTACT
7601 AGAAGATATT CTCTGGCCT CAGAAGCAGA GGTATAGAAT CTGCTCTTGT
7651 CAAGGCTTCC ACACCTCCAT AATCTACTCA ATAAACTAAG CTCAAACCTG
7701 TCCTCATTTT TATCCCCCTC ACCTCTATGT CTGATCTGTC ACAAAGTCCA
7751 GTTCATTCTT CCCATCCTTC TAATCTGTGC ACTACACTGC AACACTACCA
7801 CCGTACCCCA ACTCAGCTCA CTGCTCTCTT TTGCCTCCCA CTTTGTGAGC
7851 CTGCCTCTAG ACCTGCCCCC TCCAACCCCT CCTCCCTGTA ACCTCTACTC
7901 TTCCTAAAAC ACAATCTGAC TGAGGTATTT CCCCACTATA AACACTTCCA
7951 CAGGTGGACG AAGATGCCTC CATATCCTCT ATATGACTTC CATGCCCTTG
8001 ACAGTCTGCA CCCTGCTCCA TCTCCTCCC AGGGTCTTCC CATACCTGAT
8051 CCTCACTACA GTCCTCCTGA AATCCTGCAG GTCCTTTCAC ACATGGAGCA
8101 TCCCACACCC TTCCACATGC TCTCATCTGA CTCAAGCATC TCCTCATCTT
8151 TACTTGCA TA ATTCTACTT GTTCTTCTGG TCTTTTCATA GAGGGCCCCCT
8201 ACAGGAAGCT TCTGTGACCC CATGGCTGAG TGAGTTTCAT GTATCTATAT
8251 TGGGGTATCT GTGCATGTGT TTCCCATCA TTTGTTTAC CACATTGCAT
8301 TATAATTTTC TCTCTTCTTA CCTCACTGAT CAGACTGTGA GCTCCTGACA
8351 GGCAGGGACT TGATCGTGCC CAATTTGTCA AGGTAAACAG TGTCTGGCAC
8401 GTAGGAAGAG TTAACATCTC TGTTCTGCAT TTGTTGACTG AATGATACTG
8451 AATTTCGAAT CATCACATCA CCATGCCCT CTGTCTTAAC CATGTGAGAT
8501 ATGTGTGTGT GTGTGTATAT ATATATGTAT GTATATATAT GTATATATGT
8551 ATGTATGCAT ACATATGTAT GTATATATGT ATGTATGCAT ATATATGTAT

```

11/16

## FIGURE 3, page 4 of 9

```

8601 ATATGTATGT ATGCATATAT ATGTATATAT GTGTATATAT ACACATATAT
8651 ATGTGTGTAT ATATATAGCA AAATCACTCA AATCTTTGAG AACCATTGTC
8701 TGAAACATTC CTCATGTGAG TCTCATACAA ATATCAAGGA ACTAGCTCTG
8751 GACTCAGTGT CCCAATTTTT GTGACCAGGC ATGACCCACT CGTGTACTAA
8801 GCATGGATTA GTGGTAGTGC CTCCAATCCT CTGAAGGTCA TTTACTTCTT
8851 CTGTTGAAGG ATAACACCTA TTCTCTCCTT ACCTAACTCT AATCCTAATA
8901 GGAGCCAAAT GAGATGGTAT GCACAAAAAT GTTTTCTCAC TCTCTAGTAT
8951 TTCCCTGTGT GTAGGAATCT GGAAATTTTCTCTCAACTCA TTGAGACACA
9001 GCTGAAACTC ACCATTTCAT GGCCCATTCCT CTGAATGCCG AGGAATGCTT
9051 TGTCACTGGC TCCATGATTA CTCAGCATAT CACATCATAT TCCATGATTG
9101 ACAGTGCTGT CCTAATAATG TGTTTCTCTT CCTGCATAAC CAGGAAGAGT
9151 GGTTCCTCAT AACCACCTCTG AAGTGGGGCA CTCTCATTTG TCTTTGTGTT
9201 CTCCATGGGT ATGAGGGTTG CAGAGTCAAA AAGACTCCAT TTATTTAATG
9251 TTTTAATTCA TAACTAATGT ATCTAATGCT ATAAACTTAT AGGATATGGT
9301 TACCATTGTA TTGAAACACT CACTTGAGTA TTTTATGTCA GTCTATTTTT
9351 CTCTCCATGG CTTCCTTCCC TCTCCTTGAT TTCAGATTGT CTATGTGGCC
9401 AGAAATCCCA AGGATTGCCCT GGTGTCCTAC TACCACTTTC ACAGGATGGC
9451 TTCCTTTATG CCTGATCCTC AGAACTTAGA GGAATTTTAT GAGAAATTCA
9501 TGTCCGAAA AGGTGAGTTC AAAGTATCT TTTTGGTACC CTCTTTCAGG
9551 TGACTCTAAC AATAAGCACC TCTGTAAACT GGAGGAGAAA GTTACAAAAG
9601 GCCATCCTGA TTGAGGAGGT CCTATCTTGA TGATCTGGGA CTGGAGAAGC
9651 CAGGTAGAAG AGGTATTTTT CCAAAATTGA GTACAAATGT AAATTGAGGT
9701 CAAGTGGGGT CATAAGTTTT GAGACAAAGG TAAAAAGCCC CAAATCCTTG
9751 ACTTGAGTTT CAAACAATCA ACTTCAAAAT AAGAAGAGGC AACATTTCTT
9801 ATGGAAGAC CTGGCAGTGG GAAGGCATGA GATGATATCC TGCGGTTTCA
9851 TCTTGCAGTG TGATTGGATG GCCAGACCAC CACGTATAGT TACAAAAGTC
9901 ATACACTGCA CAGCCACAGA CAGCCTTTCC CGTAGGTCAC AGTGCACAAT
9951 CTTAGACCTG TTCACCTGCA GGAACCTCAC ATTAGAATTA GCAGCCATGA
10001 CCCCTCATCC ATTTATTAAG ATCACACCTT CCAGAGAAGC AGTGAACACA
10051 TTAGGGCCAC ACTTTTCAA ATAGCAATTG ACTAGATTG ACCAGGTGTC
10101 AAATCAGATT GGCAAGGATC TCAAACCTT CACAGAAGAA GAATATCTCA
10151 TGAACAAC AATCTCTAGA CAGAAGAAAG ATAAGATGGC TACATAAAGT
10201 GATTTAGGAT GTAAGGACCA TCTTTACATA TTTGTATGAG CATTAAATGCA
10251 GAAACATGAT TATAGTATTT CATTATAACT GCAGTGAGCA GAACCTAGAC
10301 AAATGGATAA ACATTACATG GTGTCTCCAC CATTTACTA ACGTCTCTCA
10351 AATAATTAGA ACTTCCATAA GTGAAATGGG TGAGTTGTGA TGCTGAGAGC
10401 TCCAGTCCT TGAGCAACCT AACTATCCCT GAGCAAGTGT GTTAGGGACC
10451 TGAGCTTCAT TATAACCACT CCTGACATAT TCATATTCAT ATTCTTTTGA
10501 GGAAAAAGAA AAATCAAAAT GAAGTGAAT CCACAACATG ACCTGAATAA
10551 AAACAGATAT CTGTGGAAA GTCAATAAAT AATAATTTCC ATCAAGCATG
10601 TCTCTCCAGT GAATCAAGAG AGATAACCTC ATTAGAACAT TTTCTCTAGA
10651 AACATAAATT AAAAAGGACT GACAGATGGA AATAAGAAAA ATAGCAAATC
10701 TAGATAGGAC TCCAGGTGAA TAGATTCCA ATATTTATGC AGAGAAGTTT
10751 TGAGAGCAGG AAAAGTAGGA GGAAGCAGAG GAAGACAGGA TCCAAGCTTC
10801 TTTCTTATA TGATTTTTTC CAAGGCCTAC ATTTTGTTC TGTTTTTTTG
10851 TGCAACTTG GCTGGATTGT GACACCTACA TATTTAATCA AACAACAATC
10901 CTGTTGTTTC TGTGAAAGTG TCAATGTGT AGTTGTGATT TACATCTATA
10951 ATCAGTTGAC TTTAAGAAAA GGAGATAATG TTAATATTA ATAATATGGG
11001 TGGTCTTTAT GCAATCACCT GAAGGTCTTA AGAGAAAAAA AACTCAGGTT
11051 TCCAAAGAA AGGATTCTGC CTCAGGCTG TTATATCAA TCTTGCTGTA
11101 TTTCCAGCA TCCCTACAG ATTTAAACA TGCCAAGACC CACAACCTGCA
11151 GGAGCCAATT CCTTTAATA AACCATATAT ACATAATATA CATTATATAT
11201 GTACATTTAT CTATATGTAC ATATTACATA GCCATTGCCT AAAATAAAT?
11251 ATATGTGTAC ACATTTATAT TATGCATATA TATAAACAT GCACACACAT
11301 ACATATACAT GGATGAGTCC CTTTATACT TGGTTTCCCT TTCTGCAGTT
11351 TTAGTCACCC ACAATCAACC CCAATCCAAA AATATTACAG TATTTCAAGA
11401 GAAAGAAGGA TAGAGAGAGG GAGATTACAT TCACATAGAT GTAAATATAT
11451 TAGAGAATAT TGTTATAGAA GCTCTGTTT? ATTATTAGTT TTGTTGTAAT

```

12/16

## FIGURE 3, page 5 of 9

```

11501 TCACTCACAG TGCCTAATAT AAAAATTAAA GTAATATACA CATCTATGTA
11551 TAGGAAAAAA CATAAACACT ATTATCATAT AATTTGGTAC TATCTGTGGT
11601 TTCAGGCATC CGCTGGGGGT CATGTGATAT ATCCCCTGTG CATAAGGATG
11651 AACTAATGTA TTCTTTGGAT TTGCCTATTG CTCTGTGTCT CTGGAGAACC
11701 TGGCTGACAC AGATAACAGT CACCATGACA CATGCCAAAG TTCAAGTCAC
11751 TGCAAACCTTA CATCTCTGTT TGTCTGATT CAAAAGAGAG CATTTCACAC
11801 ACTTGCTCAC TTGGTCTGTG GTCATTTTCC TCTGGAGAAT GTTTTCCATC
11851 ATCCTTCAGA CGAATCTTCA AGTCCACTTA GCACAATGCT TGCAACATAG
11901 CTCACCCCTGA ATAAAGATAG CTCCTGTGTT TATAATGACT GCCCAGAACC
11951 AAACCAGGAA GCTGCCAGAA GTTACAACCT ATCAGGGACA CTAACAATC
12001 CCTGGGATAA AATATGGTGC TGGCTAACTC AGGTGTCCAC TCATTCTCTT
12051 ACCAACTAGT GAAAAGAAAA TGCATCCCAT GTTTACCACG TAGACACAGC
12101 CTCAGCTGGA AATAGAAGTT CTCCTGGAGG GCACCCCTCT TTCTGCTGCT
12151 GCTGAGTCTC TTTTGAAGA GGAACCTGCA AACTGCATAG TGCACTATA
12201 CAGGGAAAGC AGCAGGAGGA CCCTACCCAT TTATAGGATG GCCTGAATTA
12251 GTTGAGTCTG AAACATAACA TGGTTTACCA GGAACAGGGG AAGAATTTTA
12301 TTGCCGAATG TTTTAAGACA TGTCACAAGA CATAGTCAAT GTGTGCAAAG
12351 TCACATATAA TAAATGTGTA CTATAAATCT CGGCTTTACA CCATATAGAC
12401 AAATTTTATT ACTAGAAAAT TATTTCCACT TCGTTAAGGA ACCAGAACGA
12451 TAGTTACAGA AGCTTATTTT AAAGGAGCAC TAATTTACTT TATAGCCTTG
12501 GGTTTTGTCC CAGTACTGGG AACTAACAGT GCTCTGACTT CTTCCAGTTG
12551 TTGGCGGGTC CTGGTTTGAC CATGTGAAAG GATGGTGGGC TGCAAAAGAC
12601 ATGCACCGGA TCCTCTACCT CTTCTACGAG GATATTAAAA AAGTAAGTGG
12651 CACTGAGACT TATAGGTCAG ACCCAGAAAC CCTCCTGACA ATGTTATTCT
12701 GTTAAAAAGC TGTGTCTTTA ATTGGCCAAG TTCTTCTTCT TTCCTCCCTC
12751 TTCACAATGC CTTTTTCTCC CATGATCAGA ATCCAAAACA TGAGATCCAC
12801 AAGGTGTTGG AATTCTTGGA GAAAACCTGG TCAGGTGATG TTATAAACAA
12851 GATTGTCCAC CATACTCAT TTGATGTAAT GAAGGATAAT CCCATGGCCA
12901 ACCATACTGC GGTACCTGCT CACATATTCA ATCACTCCAT CTCAAAATTT
12951 ATGAGGAAAAG GTTGGTGGCA TTTCTTTTCC TTAAGTGAAC TCTAAAAAAT
13001 TTTCTACCCT ATATGCTAAA ATAATTTTCA ACCTAATTTT CAGGCAGAAG
13051 TGAATCATTT CAGTTAAATT TTGAATCTCT GCTCCCTTCA CCCTGCCTGT
13101 TTGCAGACAG CCAATGTCAG TGGTTCTGAA ACTTGAGTCA CATTAGAACC
13151 CCTGCAGGCC TTGCTAAAGC TCTGATTGCT GGTCCCCACT CAGAGATACT
13201 GATTCCACAG ATCCAGCAGT AGCCCTCAA TTTGCTTTTC TCTCAAGTAC
13251 TCAGGTGATG CTGATGGTGC TGGTCACTTT GATTACAATA CCCACCTCAA
13301 CCATGAACCT CCCTTTGAAG GCTTGTGCAT CCTCTGAGCA GCTTTGAACA
13351 CTCATCTTTA GTCTATCCCT GTAGTTCAA ACCCTAGCTA AGCACTTAGT
13401 ACTTGGATTT GTAACACTG ATATTCATGT CTGTCTCAA AATAAGATGA
13451 TAGGCTGTCC TGAAGAGAGT GTAGTGTTCA GTTTTGTGCC ACTAGAACCT
13501 AGTATAGAGA CTCATACCTC AAAACAATC AGTAATGGCC TGTGTGTGTA
13551 GTGTACAGAT GAATGAACAT TATTTCTGTC CTCAACAAGT TAACATTCTA
13601 GATACATGCA AAAATAGCTG CAAAAAGTTA TAAACAAGAA AGTAAAGTGG
13651 AAGCTATACT AGGAATTCCC TAATACCAGT TCTCCTGGCT GTATCAAAT
13701 TACCTTTAAA AACAGTGACC CCATTCCAGA ACATTCCAAT TAACTAGTTG
13751 CAAGCTGGAA TCTAGAATTT GATATTATGG GCAAGCATTT CAGATTAATC
13801 CTCTTGTCAA AGGGTAGGAA ACCAGTAGAA ATAAAGTACT AGGATAACTT
13851 AGAGAAACAA TTCATTAGTA CAGCATTTGT TGGGCTGACA AGGTACAACA
13901 GTTTCAGAA GATCCCTAGT ATCCAAATGT CATTCCAGT GGATTACTA
13951 TTTAATTTTA CCCAACAAGT AATATCTTCT ACAATGAGGT TACTGACATC
14001 TTGTAACGTC TTTCACTGTC CCTGGGAGAA TAAGATAGGC TGTCCCCCAG
14051 GAAGTCCATG ATGGTAACCA GCTGTAGACT TTGGGTTGGG TACTAGTAGG
14101 CCAGGAGTGT ACCCTGGAGC AGAAGATTCA GCACAGTGGG GCACCTCAGT
14151 GGGGCTGCA GCATTGAGG AAGGAGTATA AGGATGCTGT GTGCCTTGTG
14201 GTAAGAAAGC CAAGTGGAGT GGAAATGAGG GACCATTTCAT AGTGGAAAGA
14251 CCAGAGGGAG TGGGACTGAG GGACCCTCAC TCATGGGAAA ATCAGTGGG
14301 TAGGCCTAGG AACCATTCAC TATAGAAAGG TTAGGTGGAA TGGGTGAGA

```

13/16

## FIGURE 3, page 6 of 9

```

14351 GTCTGTCATG AACTTCTTTG ATGTCCTACA GGGATGCCTG GAGACTGGAA
14401 GAACCACTTT ACTGTGGCTT TGAATGAGAA CTTTGATAAG CATTATGAAA
14451 AGAAGATGGC AGGGTCCACA CTGAACCTCT GCCTGGAGAT CTGAGAGGAA
14501 CAACAACAAA CTAGGTGACA GAGACTATGC CAACTATTTT GCCTTTTATT
14551 CTGTTGAGCA AGGAAGTGTG ACTGAATGTG GAGCTTATGA GCTTCAGTCC
14601 ATCTCCTATA GTGTGGCTAG TTTGCTATAA TATTAAAACA TGATTTAAAA
14651 TATCAACAAA CCAGTTACTC CAGTAAATAA AATAAGAGAA TTAGAGAGCA
14701 GAGTCCGCCT ACATGAGTTT TTTTGTTTGT TTGTTTTTTA AGTACAGGTA
14751 TGTTTTATTG TGATGACAG ACAGAGCAAA AACAAACAAT TGTCATAGG
14801 ATTTCCCATGT GTGATCCCAA GTAGATTTC CAAGAAAATT ATGCATAGGT
14851 ATTACAAGCC CCATTGTTAA GAGAGAATGT AACAGCTTGA AGTGTACATT
14901 CTATACTTTT ATGTATAAAT AATAACTTCC AAGAGAAAAG AGCTGATAAG
14951 TACATTTCAG AGTCACCATT TCTGTAATAG AATGATATAA AAATAAATTA
15001 CTAATGCAAA ATATCAATCA ATTGCAAAAT GATTACTGCT CTACTTTTGG
15051 CTTGTAACATA ATTTTCTCAA TCGAAAGAGT TTGAGTTGGA GAAATTAGTC
15101 AGTGAGTACT CCTGTAAAAA ATCTTCCCAC GATATAAATA AAATGCTTAA
15151 TGTATCTAAT CTATAAATTG AGATCTGGGA CAAATGCACC ATGACTATGC
15201 ATTGCCATTC TCTTAATAAC TATGCATTGC CATTCTCTTA ATAAAGAGTC
15251 TCTGTTGCAT CACTCTAACA ATAGGTATGA CCTCAGATTT AATATAAATT
15301 TAGTGCTTCA ACCATGCACC AGTGAAGACC TGGTTTTTTA TATGACCAAG
15351 TACAATTGTC TTATTAGAAG AGAAGGTCCT GAGGGGCCTT AGGGAGTAGG
15401 TGGATGCCAA CAGGGCTGAT GGCCTCAGAG ATGACAGCAT GTAACATGTA
15451 AGAAAGGGGG GAAATTTGGA AAGACTTAAA CTCAGAAATA AAATTAAAGA
15501 AGTGTAAGAA GGTATTCAAT TAACATCTTG AAAGGGAATC AGGAACAATA
15551 TACATAGCTC CCAGTACACA GAGCAAAGTA ACCTCCCTGT GGTGTGATC
15601 ATTGCTCTTA CCTTATCGTA GTGACAAGGC AGTGCTATTG TTACGCCAGG
15651 CAAAGGGGAA ATGACATCCT ATTCTCTAGT CAACTGGATG GAAGACAAAC
15701 TCAAAGGTAA AAAATAGATG ATGAGATTTT AAGAAAAGAG AGCAGCTGGT
15751 CCCTAAGGCC TGCTGTAAGG AAGCAGAGAG GATGAAGATG GGGGCATTG
15801 AACCAGCCCA GAGGGGACCC TGGGGTGAAA GTTCCCCTATC AGGCATTACC
15851 CCACTGCCTA CATCCAGCCA GATAACCCAA CCACCTGAAT TACCATCCTC
15901 ATATTTAGTT GGTGATCCAG AGAAGAGATT TCCTCTAGTA TTCCCTAAAG
15951 GTATAGCAAG AAAAAAGAG ATTCTTGAT CACTCCTGCC TAGTCAATGC
16001 ACTGGAAGAC TGGTTCACAG GGCTACTGTG CTTTGTAATT GGAGACATGA
16051 GAATCCACAA TGATTAGAAA GCTCGGGCCC CAGGTACTGT CAGAGTCCAC
16101 AGTCTACCTC AGAGAGGAGG AGGCAGATTA AAGGAAAAGC AAATTTTATT
16151 TTCCTACTCA GAAATGATT TCTACTAATT GAAAAGCAAT TGAATGCTGT
16201 CAATAAAGAC ATTTCTGTG CTAACCTTGG ACTCAGAGTA TTGATGACCA
16251 ACTATACAAA GCTTATTTCT TCCATGCAAT GGATGATACC TGCCTGCCTT
16301 TGCAGGTATT AAGGGGTATG ATGCTGAGAG TCCAGGTGTA GAAGATGGGA
16351 TCACGGGTTT TGGGCAAAGC ATTTTGGCTA TCTAATGTTA AGAACTGTAA
16401 GGTTTGAGAA TGCCTTGATG AAAGTTCGTA AAAGCTACAA ACAGAGTTGC
16451 TGGTCATTTT TACAAGGAGG CTGTGAACCT CTCTTCATCT TCTAGAGGCA
16501 TATTTTGGCT ATGGGCTACT AAGATTGAGA CAGGTGTAAG ATATAGTTTG
16551 CCCCATGGCC TCCCCTAGAA CTTTCCCCAA TGTGACTGTT CCTGGACTAA
16601 ACTGAGGGTC GGGCTGCTAT TTCCTGTGGC CCAATAACAA GATGCAGATG
16651 AACTGGGGAG GAAGAGAAGT TTTATTCTG TAACTGGTTA CATTGAGAGG
16701 GCCTGGAAAT TATCACCAAA CCAACTCAAA ATGACAAAAT TTTTCAGAGC
16751 TTATCTACCT TCTAAGCTGT ATGTCTACAT GTAAGTGTC ATGCCTTCTA
16801 AAGACATATG TGATTAATTT ATTTTAATTT ATAATAAGA TCTGAGTCTT
16851 GAAGACCTTC CTCTGGTGCC TAATGAAGTT TGCTTAATTT AAATGGGTCT
16901 CCAGGTACTG GGTGATCAC CCTTATCTTG TCTCCTGTTA AACTACTGAG
16951 GTTTGGGGAG TTCCTTCAGA CCTCCAATAA ACGTGTGTTG GGAGGCCTGG
17001 GGAGTTTCTT CAGAGCCCCA ATAAACTTA TATAATCCTA ACTGGGTACT
17051 GTTAAGAACT CCTTTATTAT TTTGTCATGT TGTAAGGCCC AGGAAACGCC
17101 TAGGCAAAAC TCTGGATGGG CTTTTGTTAC ATTTGAGCCT TTGCATAAGG

```

14/16

## FIGURE 3, page 7 of 9

```

17151 GCACTGGCTT TTTTAAATAT TTAACCTAAC CACTCACTGA ATACTGAAAC
17201 AGTTGTGATG GAGGCCTGCA TTAATGCAAC CTGCCTGCCA CAATCCCCAC
17251 TGTCAATTTG TGCATAATTC TTATCATGCT AGTATATTTA TTTATCATGA
17301 GAATTGTAGG GATATGGGGC ATTGTAATAT TTCTGGCTAC TTCCTGCTGA
17351 GTGAGTGTCA TTGTTATGGG ACACTGAATG CAGCATTGGT ATAGGGGAGG
17401 TCTATTTGTT CCCAGCAGCA CTCTTTGTTT CAGGGGCTTA GAGGCAGCAC
17451 CTGCTGAAAC ATTTAGTCTT CAGTTCACAG GGCTTTAAGA AAGCACAAC
17501 TAGGTTTCAG TGATTTCCAG TTAGGAAAAA TGGGGTAGTG CATGGGCTTT
17551 CATGCAGAAG AGCCTTCAGT GCAAGTCCAT GACAATGTTT GCAACTCAGT
17601 TTTATCCTCA AAAGCTCTAA CTACTTTTTC GGTATTCTAA ATCTGCTTTG
17651 CTTTAATTCG GTTTTCAAAA TATTCTTCTT CCTCAGGAAA TAGCACATGC
17701 TTTATAGTTG AATGAATCTG TTGGCTCCTG GCACTTGTA GGCAGATATC
17751 TTTATCTGCA AAATGGTATC ATATCTATCT CATGAGATCA TTGTAAAGAT
17801 TTACACTATC ATACTTGTA AGTGACATA GTTGTGTTCA AGAAATTTAG
17851 CTTCTTCTC CCTCTGCAC ACCTCAACCC CACATCATA AGCAAAGTTT
17901 AATCCATATA AACACTGGTA GGATTAGTCC CACATATTGG GACAGTGGAC
17951 TGATTTCTGA ATTTCAAACC ATCGTGCTG AGTGTCTGCA CAAATTCAC
18001 CTCATGACCC AGTGCATCAG CATCCTAGAA TTACAAAAC AGCACTGAG
18051 ATGCCTCAG AAATTTAATC AAAAGCTATA GCTTTCTCTC TTTCTGTATC
18101 TCTTTGATCA AAAAGATCCA AGAAATAAGG ATCTATGCTT CTGGGTCAAC
18151 TTGAGATTCC TTTTATTCCT CCTCTTAGTG ACCTTCTGTT AAGTTCATAG
18201 GTTCTTTTTG CCATCTGTTG TAGTCCAAAT ACTAGAGTTA GTTATACTAT
18251 GCTTCACCAC CAACATTGTT CCTCCTTCAC ACAGCCTTTC CTGTTCTCCC
18301 CCAGGAGATA TCATTGCTCC TTCCACCAA ATCTCCTGGC ACTCTGTCTG
18351 TACGTAGGCA GCATGTACTT TCTCAGCCAG CCTGTGATT GTTTGTGTTT
18401 ATGCTTAATT CCTTTCCTC AAGGTAAGCC TCTTGAGGGC AGGGACAACC
18451 CTCATCTTTG TGTTGCTACA GTATCTGGGA CAGTACAGTG TCTGGCACTA
18501 GGAGTCACTT GAGAAATATT TTTAAATGT GTTCAAATGA TCCTTTAACT
18551 CATTTCTCTC CAAATACTGT CCAATGGAA CATCATCCCC AAGGGAAGAC
18601 CCAACATGAT TTGTTTGGTG TGACTGTTCC ACATACAGGA CCCAAAGCGG
18651 GAAATTGAGA AGATACTGAA GTTCCTGGAA AAAGACATAT CAGAGGAAAT
18701 TCTGAATAAA ATCATCTATC ACACCTCCTT TGATGTAATG AAGCAAAACC
18751 CAATGACCAA CTATACCACT TTGCCACCA GCATTATGGA CCACTCCATC
18801 TCCCCTTTTA TGAGGAAAGG TAGATAAGCT TTGTAGTCTA AGATGTCAAA
18851 TGGAACCTCTG TGGTCCCCAT GGTCTGCTTA GATTTTCCAG TAATGTTTCA
18901 TTCTCCATTA TTTATTCTTT CCAGCAGCAC CACTGTACAA CCTTTGAGAG
18951 GCAAGTTGCC TGTTTCTCCT CATTCTGGT GGGGTCTTAA GGGTGCATGC
19001 TTAACCTCTC CTCTACTCCT GCAGCAATCA TTAAGATTTT GCCTTGTTTC
19051 AGGGATGCCT GGAGACTGGA AGAACTATTT TACTGTGGCC CAAATGAAG
19101 AATTGACAA GGACTACCAG AAGAAGATGG CAGGAAGCAC CCTAACCTTC
19151 CGCACAGAGA TCTGAGAGCA GTCAGGGAGT CTGTCTGGA CTTTCTTACC
19201 AGATTTTTCG CATTTGAGCC TCATGATCAA GGACACTTAA AACAAAGACA
19251 CCCTTCCTCC AGTCTGGAGC TGTTACACAC TACCTGTAGA TGATAATACT
19301 TCATCAAAAT GTAACCAAAT CCTGGGTAGA GTTTTAAAT AAGATACATG
19351 ATCCCTCATT TAGACACCAA CCACATAACG TACTCCCCTG TCCTAAGGCA
19401 AAATAAAGGC AATTAGTTC CATCACTAGT TTACAATAGT GAAATAAATA
19451 AAGAGATAAA TAAGAGTAGA ATTTCAATGT AAGAGAAAGT GAGCAAGAGA
19501 AGAATGAGGA TGATAAGTGG GCACTAAGGA TAATGTGTGG GGGAACTGGT
19551 TTTTATTTT AAAGGTCAGG AATGAGAGGC AGAGGATAGT CCTACTCTTG
19601 AAGATATTAG ATTAGGAGGT GCAGGGGGTG GCATGAAGGA TGGAATGCTG
19651 TGAGCTGGGT CTGGCTCAGG TGGCTTCAAG CTGAAATGGT CCCAAAACCA
19701 AAGAGAAAGG CCACAGGAGA GCGATAAGGA ACACAAGATG AACAAGAAAC
19751 TCAGCCTACT TCTGTGGCCA ACACAAGGCT GGAAAGCAGA GAAAGATTTT
19801 ATCTTTCAGA TGGACTTCTG AGAGAGAGAG GAAGTCTTCA GTGAGCTCAA
19851 GGGAAAATA AAATAAATA TTTGGTAACT CATGGACGTT AAGAAAAGCT
19901 TAAATTTTAC ATCTCTGTGG AAAATCGTTT TTTATACTTT TTTTAAAGG
19951 GCTCTTAGCA AAATGTCAG CATTGCCAAG TTCTTCAGCT AACTAATGA
20001 TTATGTTCTT TTCCTTCTT TTGTTAAAC CTGTACCAAG AAAATATCTG

```



15/16

## FIGURE 3, page 8 of 9

20051 CCATCATTTT ATACAAGCTA TACAACGATC CATGTATCAT TATTCTTTTA  
20101 ATGTCAGGAA GGTACATTTT GTGGGATAGG TGGTAACTT ATCATTAAAC  
20151 CATACAACAC AAAAA  
(SEQ ID NO:7)

FEATURES:Transcript #1:

Start	1009
exon	1009-1180
5'UTR	1001-1008
intron	1181-6178
exon	6179-6307
intron	6308-7158
exon	7159-7256
intron	7257-9385
exon	9386-9512
intron	9513-12547
exon	12548-12642
intron	12643-12779
exon	12880-12960
intron	12961-14381
3'UTR	14491-14514
exon	14382-14514
Stop	14491

Transcript #2:

Start	1009
exon	1009-1180
5'UTR	1001-1008
intron	1181-6178
exon	6179-6307
intron	6308-7158
exon	7159-7256
intron	7257-9385
exon	9386-9512
intron	9513-12547
exon	12548-12642
intron	12643-18638
exon	18639-18819
intron	18820-19052
exon	19053-19165
Stop	19163

cDNA:

exon	1011-1180
5'UTR	1001-1008
intron	1181-6178
Start	1009
exon	6179-6307
intron	6308-7158
exon	7159-7256
intron	7257-9385
exon	9386-9512
intron	9513-12547
exon	12548-12642
intron	12643-12779
exon	12880-12960

16/16

## FIGURE 3, page 9 of 9

intron 12961-14381  
 exon 14382-14506  
 Stop 14491  
 3'UTR 14491-14506

Map position:

H1552	#	SHGCNAME	CHROM#	LOD_SCORE	DIST. (cRs)
	1	SHGC-1639	2	10.01	16

Variants/SNPs:

Position	Major	Minor	Protein coding/non-coding				
1,008	a	c	exon, outside orf				
6,094	a	g	intron				
7,898	t	c	intron				
10,018	a	c	intron				
10,791	a	t	intron				
12,451	a	t	intron				
14,674	t	c	intron				
14,860	g	c	intron				
20,133	a	g	intron				
1,025	a	c	exon	39	V	V	Protein 3
12,602	t	c	exon	194	M	T	Protein 3
12,829	g	t	exon	224	W	L	Protein 3
14,420	a	t	exon	281	L	M	Protein 3

Context of variation:DNA

Position	
1,008	ttagtattgatcttaccatcccagattccc[a/c]atggcgaagattgagaaaaa (SEQ ID NO: 11)
6,094	gcaacattgatctaattttaaagtgtctctcat[a/g]ttgctgtctttcttcaaaatattc (SEQ ID NO: 12)
7,898	ccaacccctcctcctgtaacctcta[t/c]tcttcctaaaacacaatct (SEQ ID NO: 13)
10,018	agcagccatgaccctcatccattttatt[a/c]agatcacaccttccagagaa (SEQ ID NO: 14)
10,791	aggaggaagcagaggaagacagga[a/t]ccaagcttctttcttatatgattttttcc (SEQ ID NO: 15)
12,451	ccacttcgttaaggaaccagaacga[a/t]agttacagaagcttatttcaaaggag (SEQ ID NO: 16)
14,674	tttaaatatcaacaaccagttactccag[t/c]aaataaataagagaattagagagcagagtc (SEQ ID NO: 17)
14,860	aggtattacaagc[g/c]ccattgttaagagagaatgtaacagctt (SEQ ID NO: 18)
20,133	ttttgtgggataggtg[a/g]ttaacttatcattaaacca (SEQ ID NO: 19)
1,025	GGTGGGAAAAAGT[a/c]TGTAATTTCCAAG (SEQ ID NO: 20)
12,602	GTGGGCTGCAAAAGACA[t/c]GCACCGGATC (SEQ ID NO: 21)
12,829	GGAGAAAATT[g/T]GTCAGGTGATGTT (SEQ ID NO: 22)
14,420	ACTTTACTGTGGCT[A/t]TGAAT (SEQ ID NO: 23)

## SEQUENCE LISTING

<110> Woodage, Trevor  
Wei, Minh Hui  
Kodira, Chinappa  
Beasley, Ellen  
DiFrancesco, Valentina

<120> ISOLATED HUMAN DRUG-METABOLIZING  
PROTEINS, NUCLEIC ACID MOLECULES ENCODING HUMAN  
DRUG-METABOLIZING PROTEINS, AND USES THEREOF

<130> CL000669PCT

<140> N/A

<141> 2001-0327

<150> 60/192,408

<151> 2000-03-27

<150> 60/212,725

<151> 2000-06-20

<150> 09/609,816

<151> 2000-07-03

<160> 23

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 942

<212> DNA

<213> HUMAN

<400> 1

```
attcccaatg gcgaagattg agaaaaacgc tcccacgatg gaaaaaaagc cagaactgtt 60
taacatcatg gaagtagatg gagtcacctac gttgatatta tcaaaagaat ggtgggaaaa 120
agtatgtaat ttccaagcca agcctgatga tcttattctg gcaacttacc caaagtcagg 180
tacaacatgg atgcatgaaa ttttagacat gattctaaat gatggtgatg tggagaaatg 240
caaaagagcc cagactctag atagacacgc ttcccttgaa ctgaaatttc cccataaaga 300
aaaaccagat ttggagtctg ttcttgaaat gtcctcacca caactgataa aaacacatct 360
cccttcacat ctgattccac catctatctg gaaagaaaac tgcaagattg tctatgtggc 420
cagaaatccc aaggattgcc tgggtgtccta ctaccacttt cacaggatgg cttectttat 480
gcctgatacct cagaacttag aggaatttta tgagaaattc atgtccgga aagttgttgg 540
cggggtcctgg tttgaccatg tgaaaggatg gtgggctgca aaagacatgc accggtatcct 600
ctacctcttc tacgaggata ttaaaaaaaa tccaaaacat gagatccaca aggtgttggg 660
attcttgagg aaaacttggc cagggtgatgt tataaacaag attgtccacc atacctcatt 720
tgatgtaatg aaggataatc ccatggccaa ccatactgcg gtacctgctc acatattcaa 780
tactccatc tcaaaattta tgaggaaagg gatgcctgga gactggaaga accactttac 840
tgtggctttg aatgagaact ttgataagca ttatgaaaag aagatggcag ggtccacact 900
gaacttctgc ctggagatct gagaggaaca acaacaaact ag 942
```

<210> 2

<211> 922

<212> DNA

<213> HUMAN

<400> 2

```
attcccaatg gcgaagattg agaaaaacgc tcccacgatg gaaaaaaagc cagaactgtt 60
taacatcatg gaagtagatg gagtcacctac gttgatatta tcaaaagaat ggtgggaaaa 120
agtatgtaat ttccaagcca agcctgatga tcttattctg gcaacttacc caaagtcagg 180
tacaacatgg atgcatgaaa ttttagacat gattctaaat gatggtgatg tggagaaatg 240
```

```

caaaagagcc cagactctag atagacacgc ttctcttgaa ctgaaatttc cccataaaga 300
aaaaccagat ttggagttcg ttcttgaaat gtctcacca caactgataa aaacacatct 360
cccttcacat ctgattccac catctatctg gaaagaaaac tgcaagattg tctatgtggc 420
cagaaatccc aaggattgccc tgggtgccta ctaccacttt cacaggatgg cttcctttat 480
gcctgatcct cagaacttag aggaatttta tgagaaattc atgtccggaa aagttgttgg 540
cgggtcctgg ttgaccatg tgaaaggatg gtgggctgca aaagacatgc accggatcct 600
ctacctcttc tacgaggata ttaaaaaaga cccaaagcgg gaaattgaga agatactgaa 660
gttctctggaa aaagacatat cagaggaaat tctgaataaa atcatctatc acacctcctt 720
tgatgtaatg aagcaaaacc caatgaccaa ctataccact ttgccacca gcattatgga 780
ccactccatc tcccctttta tgaggaaagg gatgcctgga gactggaaga actattttac 840
tgtggcccaa aatgaagaat ttgacaagga ctaccagaag aagatggcag gaagcaccct 900
aaccttccgc acagagatct ga                                     922

```

&lt;210&gt; 3

&lt;211&gt; 1004

&lt;212&gt; DNA

&lt;213&gt; HUMAN

&lt;400&gt; 3

```

ctcactatta gggcgaattg aatttagcgg ccgcgaattc gcccttatgg cgaagattga 60
gaaaaacgct cccacgatgg aaaaaagcc agaactgttt aacatcatgg aagtagatgg 120
agtccttacg ttgatattat caaaagaatg gtgggaaaaa gtctgtaatt tccaagccaa 180
gcctgatgat cttattctgg caacttacct aaagtcaggt acaacatgga tgcataaat 240
tttagacatg attctaaatg atggtgatgt ggagaaatgc aaaagagccc agactctaga 300
tagacacgct ttcttgaaac tgaaatttcc ccataaagaa aaaccagatt tggagtctgt 360
tcttgaaatg tctcaccac aactgataaa aacacatctc ccttcacatc tgattccacc 420
atctatctgg aaagaaaact gcaagattgt ctatgtggcc agaaatccca aggattgcct 480
ggtgtcctac taccactttc acaggatggc ttcttttatg cctgatcctc agaacttaga 540
ggaattttat gagaaattca tgtccggaaa agttgttggc gggtcctggt ttgaccatgt 600
gaaaggatgg tgggctgcaa aagacacgca ccggtcctc tacctcttct acgaggatat 660
taaaaaaaat ccaaaacatg agatccacaa ggtgttgaa ttcttgagaa aaactttgtc 720
aggatgatgt ataaacaaga ttgtccacca tacctcattt gatgtaatga aggataatcc 780
catggccaac catactgcgg tacctgctca catattcaat cactccatct caaaatttat 840
gaggaaaggg atgcctggag actggaagaa ccactttact gtggctatga atgagaactt 900
tgataagcat tatgaaaaga agatggcagg gtccacactg aacttctgcc tggagatctg 960
agaggaacaa caaagggcga attcgtttaa acctgcagga ctag                                     1004

```

&lt;210&gt; 4

&lt;211&gt; 304

&lt;212&gt; PRT

&lt;213&gt; HUMAN

&lt;400&gt; 4

```

Met Ala Lys Ile Glu Lys Asn Ala Pro Thr Met Glu Lys Lys Pro Glu
 1           5           10           15
Leu Phe Asn Ile Met Glu Val Asp Gly Val Pro Thr Leu Ile Leu Ser
          20           25           30
Lys Glu Trp Trp Glu Lys Val Cys Asn Phe Gln Ala Lys Pro Asp Asp
          35           40           45
Leu Ile Leu Ala Thr Tyr Pro Lys Ser Gly Thr Thr Trp Met His Glu
          50           55           60
Ile Leu Asp Met Ile Leu Asn Asp Gly Asp Val Glu Lys Cys Lys Arg
          65           70           75           80
Ala Gln Thr Leu Asp Arg His Ala Phe Leu Glu Leu Lys Phe Pro His
          85           90           95
Lys Glu Lys Pro Asp Leu Glu Phe Val Leu Glu Met Ser Ser Pro Gln
          100          105          110
Leu Ile Lys Thr His Leu Pro Ser His Leu Ile Pro Pro Ser Ile Trp
          115          120          125
Lys Glu Asn Cys Lys Ile Val Tyr Val Ala Arg Asn Pro Lys Asp Cys
          130          135          140
Leu Val Ser Tyr Tyr His Phe His Arg Met Ala Ser Phe Met Pro Asp
          145          150          155          160

```

Pro Gln Asn Leu Glu Glu Phe Tyr Glu Lys Phe Met Ser Gly Lys Val  
 165 170 175  
 Val Gly Gly Ser Trp Phe Asp His Val Lys Gly Trp Trp Ala Ala Lys  
 180 185 190  
 Asp Met His Arg Ile Leu Tyr Leu Phe Tyr Glu Asp Ile Lys Lys Asn  
 195 200 205  
 Pro Lys His Glu Ile His Lys Val Leu Glu Phe Leu Glu Lys Thr Trp  
 210 215 220  
 Ser Gly Asp Val Ile Asn Lys Ile Val His His Thr Ser Phe Asp Val  
 225 230 235 240  
 Met Lys Asp Asn Pro Met Ala Asn His Thr Ala Val Pro Ala His Ile  
 245 250 255  
 Phe Asn His Ser Ile Ser Lys Phe Met Arg Lys Gly Met Pro Gly Asp  
 260 265 270  
 Trp Lys Asn His Phe Thr Val Ala Leu Asn Glu Asn Phe Asp Lys His  
 275 280 285  
 Tyr Glu Lys Lys Met Ala Gly Ser Thr Leu Asn Phe Cys Leu Glu Ile  
 290 295 300

&lt;210&gt; 5

&lt;211&gt; 304

&lt;212&gt; PRT

&lt;213&gt; HUMAN

&lt;400&gt; 5

Met Ala Lys Ile Glu Lys Asn Ala Pro Thr Met Glu Lys Lys Pro Glu  
 1 5 10 15  
 Leu Phe Asn Ile Met Glu Val Asp Gly Val Pro Thr Leu Ile Leu Ser  
 20 25 30  
 Lys Glu Trp Trp Glu Lys Val Cys Asn Phe Gln Ala Lys Pro Asp Asp  
 35 40 45  
 Leu Ile Leu Ala Thr Tyr Pro Lys Ser Gly Thr Thr Trp Met His Glu  
 50 55 60  
 Ile Leu Asp Met Ile Leu Asn Asp Gly Asp Val Glu Lys Cys Lys Arg  
 65 70 75 80  
 Ala Gln Thr Leu Asp Arg His Ala Phe Leu Glu Leu Lys Phe Pro His  
 85 90 95  
 Lys Glu Lys Pro Asp Leu Glu Phe Val Leu Glu Met Ser Ser Pro Gln  
 100 105 110  
 Leu Ile Lys Thr His Leu Pro Ser His Leu Ile Pro Pro Ser Ile Trp  
 115 120 125  
 Lys Glu Asn Cys Lys Ile Val Tyr Val Ala Arg Asn Pro Lys Asp Cys  
 130 135 140  
 Leu Val Ser Tyr Tyr His Phe His Arg Met Ala Ser Phe Met Pro Asp  
 145 150 155 160  
 Pro Gln Asn Leu Glu Glu Phe Tyr Glu Lys Phe Met Ser Gly Lys Val  
 165 170 175  
 Val Gly Gly Ser Trp Phe Asp His Val Lys Gly Trp Trp Ala Ala Lys  
 180 185 190  
 Asp Met His Arg Ile Leu Tyr Leu Phe Tyr Glu Asp Ile Lys Lys Asp  
 195 200 205  
 Pro Lys Arg Glu Ile Glu Lys Ile Leu Lys Phe Leu Glu Lys Asp Ile  
 210 215 220  
 Ser Glu Glu Ile Leu Asn Lys Ile Ile Tyr His Thr Ser Phe Asp Val  
 225 230 235 240  
 Met Lys Gln Asn Pro Met Thr Asn Tyr Thr Thr Leu Pro Thr Ser Ile  
 245 250 255  
 Met Asp His Ser Ile Ser Pro Phe Met Arg Lys Gly Met Pro Gly Asp  
 260 265 270  
 Trp Lys Asn Tyr Phe Thr Val Ala Gln Asn Glu Glu Phe Asp Lys Asp  
 275 280 285

Tyr Gln Lys Lys Met Ala Gly Ser Thr Leu Thr Phe Arg Thr Glu Ile  
 290 295 300

<210> 6  
 <211> 304  
 <212> PRT  
 <213> HUMAN

<400> 6  
 Met Ala Lys Ile Glu Lys Asn Ala Pro Thr Met Glu Lys Lys Pro Glu  
 1 5 10 15  
 Leu Phe Asn Ile Met Glu Val Asp Gly Val Pro Thr Leu Ile Leu Ser  
 20 25 30  
 Lys Glu Trp Trp Glu Lys Val Cys Asn Phe Gln Ala Lys Pro Asp Asp  
 35 40 45  
 Leu Ile Leu Ala Thr Tyr Pro Lys Ser Gly Thr Thr Trp Met His Glu  
 50 55 60  
 Ile Leu Asp Met Ile Leu Asn Asp Gly Asp Val Glu Lys Cys Lys Arg  
 65 70 75 80  
 Ala Gln Thr Leu Asp Arg His Ala Phe Leu Glu Leu Lys Phe Pro His  
 85 90 95  
 Lys Glu Lys Pro Asp Leu Glu Phe Val Leu Glu Met Ser Ser Pro Gln  
 100 105 110  
 Leu Ile Lys Thr His Leu Pro Ser His Leu Ile Pro Pro Ser Ile Trp  
 115 120 125  
 Lys Glu Asn Cys Lys Ile Val Tyr Val Ala Arg Asn Pro Lys Asp Cys  
 130 135 140  
 Leu Val Ser Tyr Tyr His Phe His Arg Met Ala Ser Phe Met Pro Asp  
 145 150 155 160  
 Pro Gln Asn Leu Glu Phe Tyr Glu Lys Phe Met Ser Gly Lys Val  
 165 170 175  
 Val Gly Gly Ser Trp Phe Asp His Val Lys Gly Trp Trp Ala Ala Lys  
 180 185 190  
 Asp Thr His Arg Ile Leu Tyr Leu Phe Tyr Glu Asp Ile Lys Lys Asn  
 195 200 205  
 Pro Lys His Glu Ile His Lys Val Leu Glu Phe Leu Glu Lys Thr Leu  
 210 215 220  
 Ser Gly Asp Val Ile Asn Lys Ile Val His His Thr Ser Phe Asp Val  
 225 230 235 240  
 Met Lys Asp Asn Pro Met Ala Asn His Thr Ala Val Pro Ala His Ile  
 245 250 255  
 Phe Asn His Ser Ile Ser Lys Phe Met Arg Lys Gly Met Pro Gly Asp  
 260 265 270  
 Trp Lys Asn His Phe Thr Val Ala Met Asn Glu Asn Phe Asp Lys His  
 275 280 285  
 Tyr Glu Lys Lys Met Ala Gly Ser Thr Leu Asn Phe Cys Leu Glu Ile  
 290 295 300

<210> 7  
 <211> 20165  
 <212> DNA  
 <213> HUMAN

<400> 7  
 agttccaaac tttccacat tttcctgtct tcttctgttc caatctctgc ctgttaccca 60  
 gttccaaaagt caattttaca ttttcaggtta tctgcagtag caccocactc tgctagtaca 120  
 aattttattgt attagtcctt tttcatgctg ttaataaaga catacctgag actgggcaat 180  
 ttaaaaaata acgaggttta acagacttaa agttccacgt atctggggaa gcctcacagt 240  
 catggtggaa ggcaatgaat ggcagcaggc aaaggaaaag agagcttgtg caggggaact 300  
 ccccttata atgggtggg acacaggcga accatatcag accccaacta aattccaagt 360  
 ttccagagtt agtagacatg agagttttca ggtacatggg tacaagagag ttttttctgc 420

tgtaacccat gactgatatt tcaaaaatta ttccatgaat gaaaaaaaaa ctacatgaaa 480  
ttatgttttt tgaagatttt ctctgtgaaa actattcaag aaaattgagt atagaatgct 540  
ccttaaaacc attgttttga attttcttca atgtaattgt ctgccttcta attatacaaa 600  
ataaatatctt gaagacaatg aagcaatata tgacacataa cccattatat tcccagattaa 660  
caagtaatat ggttgtagg tcatgcagca ggaaaaagac tatttgggta taaatctcag 720  
ccccaacagg tattgattat ttacttttta tcaaattatt taaacttata atgacttaag 780  
tttccacatg tagaaaacag aaaaaactat tacctatttt atgtgggtgc tatgaagatt 840  
aaataattaa tgtacataga gtaagttgtt agcatgttat atgttagctt tcacttattt 900  
tatgtctatt taatcaaagg cagtaagatg ataaccata ctatgaacca taaaacaaca 960  
tttttaaaaa ttttaattag tattgatctt acccatccca gattcccaat ggcgaagatt 1020  
gagaaaaacg ctcccacgat ggaaaaaaag ccagaactgt ttaacatcat ggaagtagat 1080  
ggagtcccta cgttgatatt atcaaaagaa tgggtgggaaa aagtatgtaa tttccaagcc 1140  
aagcctgatg atcttattct ggcaacttac ccaaagtcag gtaagggtag caaaacataa 1200  
aaatattcaa tattttcacg tgaattatt gcataatctg tattgataaa tgaagcatga 1260  
ttgggatttg gagagaaaaca attcctcatt atggagatct gttctttggt gctgcaggac 1320  
atttagctt cctgatccct aggacaacaa ttccaatag cactctctga agagaacaga 1380  
aaagattttct aatgctttg gtagtagggc agtaaatgct cccattccag aacctatagc 1440  
atacatcaaa tataagggtca acaaattgaa agacctatgg cttttttaaa catcaaccct 1500  
caagacagcc ttctaaaagc atgtcctact ccaaattatta atctattatc tcagatatta 1560  
aacacaaatt gattttctaa tcctctaaag ctcttgagg atgtcagcca tgaattttct 1620  
gctctgtacc aaattagtct cattcagaaa agcccaatga ctgacctga tttaaaatgc 1680  
cttagatttc taattcttct ctaaaaattc ctgactgga acacatgcta gagtcaatgg 1740  
gcacacctaa tgcccagaac tcactttcga taggccattc tccattaaaa tgaaccacag 1800  
ctattaggag aattgatgat cccactcttg ggaatagaat atgcaagggg aatgtactac 1860  
atcttctttt tgaaggagta agtgatcagt atatatctcc agaattgttt gtttgtatta 1920  
aatgtctgta aaaataaagc acacataata acaaaaaaat tgatggggac atgtagaagg 1980  
atacaagaac caacgtgaag gggctctcac agtagccaca tttgggacag tttgagcatc 2040  
caaaagaatg atcactaact gattgggaga acattaaata aaaacctcct ggtcagccat 2100  
gccgaaggga caagattgga tgtgtactcc agccataata aaaaaggaaa ctggataaaa 2160  
tatatgaaac cactggttc tgacctgga tgacagttgt ctcaatcctc agactgaaga 2220  
tgctttctga atagggtata gttatagggt gcaaaagaaa aagccatagc aaactcattg 2280  
agttcaagag ccagggtgtag gagttctgag aggttgaagt ggctgctatt tgcaggtaag 2340  
agtaccagac agaaggaaac tacacaaata aataccttta gatattctta gagaggaccc 2400  
ttttagttta ttgttgaatg gtagactgca cttccataga gcctatagct ccatgagatc 2460  
aggcaaggaa ccaccagaaa actattagcc atataattcc tagagattat acaagcatga 2520  
gagacttttg tgctccaatc aggataggat ggacagaatt ttggtcccca tgacattagt 2580  
cctctggtat tacatctgcc gttatttcag attacattcc caaaaggatt ttgcagttgc 2640  
tattaaacta tctaatacgc tgatattaaa ataggagat tattctggac tatcttgacg 2700  
gacccagtgg aatcatgtga acaggaagtc agagagatgt ggcagaggag aaagtctgag 2760  
aaattttaa gataataaaa gttcactgca tgattataga tttaatgatg aagagagaaa 2820  
gtatcaacaa aacagttatc tcaatactac agccacaagc aactgaattc tgttggcatc 2880  
taggagcttg gcaaaacacc ctgaagtccc agatgagaat ggcagcccta gctgatacct 2940  
tgatttttag ctagtgagac cctaacaga ggactagcca tgttaacccc aatttctaatt 3000  
ctacagaaac tatgaccgaa tattcagggt ttgttttaac tcatgaagct tgtggtaatt 3060  
tgtaaacac aaagtcttca acctagaatt ccatataccc agtaaaaaata atatatataa 3120  
atattactaa aaataaagg caaagacttt ttctctgaca gcgaaaagct gaatgtgttg 3180  
ccgcacagc tgcactaaaa ataaataaat aaaaagttat attaaagaaa tttattcaga 3240  
gtaacagaaa ataatagtat acaaaaacttt attatctttc aacattccag caacacacag 3300  
ttggaatatg acattttaaa aataccattg ataaaaataa tagcacctga aaacatgaaa 3360  
tattcagaaa tgaacttaac aaagctgtgc atgatttgaa agtgaaaaca ctacggagag 3420  
aaattgtata agacttaaat aaataaggag gcatacccta tctacggatc aaaaggatca 3480  
atatttttaa gatatcaact gtttcaaaat taatgtataa attcaatctc aatcacaac 3540  
tgacaaattg attccaagat ttctatgaaa atgtcattaa aagcaaatat tttgataga 3600  
gtaaaagtggc aggggtttata ttaactgatt tcatgacatt taactcagct tcaacattca 3660  
aaagactgtg atactgttga tattgttgat agacataaat acttgacct tacttgggtg 3720  
cagatgcaaa aaagtcaaag tgcaatgcat cagacctcaa aaaattaaaa ttaaatctct 3780  
aaaactatgg aagtatatag aggagaatag ttccaactt taggttaggc aacaataatt 3840  
tggggaaaaa aatggaaagc actacccttt atggtttgca tttaatgtaa attcaatata 3900  
aattagactt aatcaaatga aaacttctgt tcttctaaag acccagttaa gaaaaataaa 3960  
atatgtgaca tagatggaga gaaaatattc acaatacata tatctggcca gaaggataaa 4020  
agaactgtta caactaagaa caaaaaacaa aaaaaaatg tattaaagtg ggcaaaagat 4080  
gtaaaaatgt atcaccaaaa aagctttact attaaccaat aaacacaaga tattcaacat 4140  
catttatcat gagaaaatgt aaattagtac tataatgaaa taccactaca caccatttca 4200

aatggctaaa	aacctggaaa	tactaagtgt	tgataaggac	acagagcaac	tgaaattctc	4260
atgaactttt	ggggaggacc	taacatggta	ctaagtgtga	acagtttgtt	ataaaactaa	4320
acatgcacct	accacacaac	ccagcaactc	caatcctaga	gattcaccce	cacccccaaa	4380
atagaaatct	atgttcatat	aaaaacttgt	acacaacact	atagcagact	tattcaaaat	4440
tagcccaaca	ctgaagaaaa	cctaaattcg	tattgtcata	taaatgaata	aacaaatcac	4500
tgcttatcca	tacaattgaa	ctgttctcag	caattaaaaa	aaatgaacta	cagatatata	4560
caagaacata	aatgaatccc	aaaataatga	gggtgtgtga	aaaaattcag	acacatgagg	4620
acaaactgta	tggtttcatt	tatatgaaat	tcaacaatgt	gtaaaactaa	tttgatttga	4680
caaaaacata	gatcagcagt	tccgtggagt	cgggaagtga	gaaaagaatt	actgatagca	4740
acagacacaa	cgttctactc	ttatctatgg	taatggctac	atacatttgt	caaatttctc	4800
taaactgacac	attttataaca	ggttcattct	attctatgta	atttttacc	cactgaagtt	4860
tatttcaaaa	aacgtgattc	cataatgatg	aaaaatacaa	aaagaaaaac	ttatgtatta	4920
tgattgaaga	taactgttat	tcatctctta	gactaaaaag	aagtaattaa	gagaaagaat	4980
ttagaggaat	tgacgttctt	ccctgattta	tgagggaag	ttctttatat	gaagatctac	5040
ctaataaata	gagaagtga	gggattagaa	aataagcaat	ttgaaaccca	caataaaaa	5100
taggtaaagt	aggatcatta	atgaatgaat	cttaaaaata	ttagataaaa	tatgaggaga	5160
actggctggg	cacatggtag	tgaaatgtca	ccacatagtt	ttattcctaa	aagcaaatgg	5220
aaatatgggc	cttttacaatg	agcgacctgg	tggtcaccta	caaaaactga	tgatgaacag	5280
tagtatcact	agtagtgagg	caatcagacg	tatgtacttc	ttggcatggg	gcaagaagta	5340
ccaagcactg	cctgagaaat	ctgccaaaaa	tggtcgctag	aacctaatca	agacctggat	5400
tccagtttat	gggaaataga	ggaaatagag	gaacaaaaat	gtactgtgaa	gaaataatca	5460
gacaaatcca	gaatgtgggg	agtactatag	gacagctggc	taattcgttc	aaaaaccaat	5520
gtcaaagaac	aaaaatcagt	atcattaaga	aaaaatatgg	aaagatgatt	ctaagctgag	5580
acaaaaagga	tagcaaaagt	aatgcagcca	atcaccttta	attgggcaaa	agtgattaat	5640
aacattatga	tacaatggaa	aaaaattaac	ataggttgag	tatctcatgt	tattaaggat	5700
ttattataaa	ctagatgtga	tgagaaacat	ttagtaacca	tgataagaga	atgtcatatt	5760
ggaagataca	gcatttttgg	catgaagtgt	cataatatag	gaaatgtaga	aatgggtcac	5820
tcaaactatga	tagttgagag	attaaatact	aaaagatgat	agatagatag	acagataata	5880
ggaagatagt	agatacatat	atacatatcat	acttagatag	atagatagat	acatagatag	5940
ataaagaana	agatgataga	ttagatagat	agatagatag	atagatagat	agatagatag	6000
atggatagat	agatgagtag	atagatgcag	gtataatata	acagttttca	tttctgcctt	6060
ggcaacattg	atctaatttt	aaagtgtctc	catgttgctg	tctttcttca	aaatattcaa	6120
taacttagaa	tgaaggttca	attgactaaa	attaaagaac	tatttcaaat	attttcaggt	6180
acaacatgga	tgcatgaaat	tttagacatg	attctaaatg	atgggtgatg	ggagaaatgc	6240
aaaagagccc	agactctaga	tagacacgct	ttccttgaac	tgaaatttcc	ccataaagaa	6300
aaaccaggtg	agtaatatgc	acgaagatag	aaaggacttt	cacttcagga	ttccagagca	6360
atgtgtactg	tctctgatag	agcatccgtg	gtagccagaa	gtagcctgta	tctttcatga	6420
ggcttatttg	ttctcagggc	aacaatcaaa	ctctagattg	ggcttccagg	gcttctcctt	6480
gttcttagct	gggtggcctt	atctcccaaa	taagattttc	attctctttc	tcatatttct	6540
cctacccaat	gttacgataa	agaagactcc	ctctgctttg	tatttctcct	catatgtttg	6600
aatcagtgga	aaggccagaa	attagcaatc	agctattgta	aaataacagg	tgtgatttct	6660
gaagaaaaag	gaggagaagt	aggggggata	taccgtggac	attcccagaa	tttaattggg	6720
gagctgaaaa	ggtttctctc	gagtgtaaaa	cccagtgaat	aacatgaaat	aaagacagac	6780
ctcctagagag	tctttctttc	caagaatgac	aacctattgt	aaaaaaatgg	atgtaatttc	6840
tgacaaacaa	aaggaagaga	gaaaaagaga	aaggaaaaaa	acaaccagat	atataacaaa	6900
cattcccaga	attaatcgag	gggtgaaaaa	agctttcctg	aaagtctaaa	agctgagaga	6960
ataagaggaa	ataagcaaaa	ctctcctaga	aggttaggca	ggattggatg	ctttgtatgt	7020
ccttgtggat	aaaaaattac	tatttccaat	ggatataaaa	caaaaataaa	tactcttaca	7080
gacaaagata	taaatggaat	tcaagtattt	tggcagagtg	ccaagaataa	acaaagaata	7140
taaattgttt	cttgctagat	ttggagtctg	ttcttgaaat	gtcctcacca	caactgataa	7200
aaacacatct	ccttccacat	ctgattccac	catctatctg	gaaagaaaac	tgcaaggtag	7260
aaagaggggg	cttttcaaac	ttctcttagc	ttggtgatag	aaactataca	actgaagata	7320
tctttcaaaa	taataatact	tgaaaaatat	tttccaaaat	ataatttgct	atttttctta	7380
gatgaagcac	ttaaaatcaa	ggattacata	aatttgaaat	ctgcaaacat	ccatgttttc	7440
taaaattcat	tttctcttaa	tcctatttca	tgaaaaattc	ttggttaagat	tttccaaaat	7500
tgagtctgtg	ttgctacaaa	tcagagagtg	tattggggag	tagaatggag	ggaacacat	7560
ttaaaaataa	aacctgtttg	ctttgctctg	ccaaagtact	agaagatatt	ctcctggcct	7620
cagaagcaga	ggtatagaat	ctgctcttgc	caaggcttcc	acacctccat	aatctactca	7680
ataaaactag	ctcaaacctg	tcctcatatt	tatccccctc	acctctatgt	ctgatctgtc	7740
acaaagtcca	gttcattctt	cccatccttc	taatctgtgc	actacactgc	aacactacca	7800
cctgtaccca	actcagctca	ctgctctctt	ttgcctccca	ctttgtcagc	ctgcctctag	7860
acctgcccc	tccaacccct	cctccctgta	acctctactc	ttcctaaaac	acaatctgac	7920
tgaggtagtt	ccccactata	aacacttcca	cagggtggag	aagatgcctc	catatcctct	7980



atatgacttc	catgcccttg	acagtctgca	ccctgctcca	tctcctcccc	agggtcttcc	8040
catacctgat	cctcactaca	gtcctcctga	aatcctgcag	gtcctttcac	acatggagca	8100
tccccacacc	ttccacatgc	tctcatctga	ctcaagcacc	tctcatctt	tacttgata	8160
attcctactt	gttcttctgg	tcttttcata	gagggccct	acaggaagct	tctgtgacct	8220
catggctgag	tgagtttcat	gtatctatat	tgggttatct	gtgcatgtgt	ttccccatca	8280
ttttgttcac	cacattgcat	tataattttc	tctcttctta	cctcactgat	cagactgtga	8340
gctcctgaca	ggcagggact	tgatcgtgcc	caatttgtca	aggtaaacag	tgtctggcac	8400
gtaggaaag	ttaacatctc	tggtctgcat	ttgttgactg	aatgatactg	aatttccaat	8460
catcacatca	ccatgcccct	ctgtcctaac	catgtgagat	atgtgtgtgt	gtgtgtatat	8520
atatatgtat	gtatatatat	gtatatatgt	atgtatgcat	acatatgtat	gtatatatgt	8580
atgtatgcat	gtatatgtat	atgtatgtat	atgtatatat	atgtatatat	gtgtatatat	8640
acacatatat	atgtgtgtat	atatatagca	aaatcactca	aatctttgag	aaccattgtc	8700
tgaacatttc	ctcatgtgag	tctcatacaa	atatcaagga	actagctctg	gactcagtgt	8760
cccaattttt	gtgaccaggc	atgaccact	cgtgtactaa	gcatggatta	gtggtagtgc	8820
ctccaatcct	ctgaagggtca	tttacttctt	ctgttgaagg	ataacaccta	ttctctcctt	8880
acctaactct	aatcctaata	ggagccaaat	gagatggtat	gcacaaaaat	gttttctcac	8940
tctctagtat	ttccccgtgt	gtaggaatct	ggaaaatttc	tctcaactca	ttgagacaca	9000
gctgaaactc	accatttcat	ggcccccttc	ctgaatgccc	aggaatgctt	tgtcactggc	9060
tccatgatta	ctcagcatat	cacatcatat	tccatgattg	acagtgtgtg	cctaataatg	9120
tggtcatctt	cctgcataac	caggaagagt	ggttcctcat	aaccactctg	aagtggggca	9180
ctctcatttg	tctttgtgtt	ctccatgggt	atgaagggtg	cagagtcaaa	aagactccat	9240
ttatttaatg	ttttaattca	taactaatgt	atctaattgt	ataaacttat	aggatatggt	9300
taccattgta	ttgaaacact	cacttgagta	tttcatgtca	gtctattttt	ctctccatgg	9360
cttccttccc	tctccttgat	ttcagattgt	ctatgtggcc	agaaatccca	aggattgcct	9420
gggtgcctac	taccactttc	acaggatggc	ttcctttatg	cctgatcctc	agaacttaga	9480
ggaattttat	gagaaattca	tgtccggaaa	aggtaggttc	aaactgatct	ttttggtacc	9540
ctctttcagg	tgactctaac	aataagcacc	tctgtaaact	ggaggagaaa	gttacaaaag	9600
gccatcctga	ttgaggaggt	cctatcttga	tgatctggga	ctggagaaag	caggtagaag	9660
aggatttttt	ccaaaattga	gtacaaatgt	aaattgaggt	caactggggt	cataagtttt	9720
gagacaaagg	taaaaagccc	caaatccttg	acttgagttt	caaacaatca	acttcaaaat	9780
aagaagaggc	aacatttcct	atggaaagac	ctggcagtgg	gaaggcatga	gatgatattc	9840
tgcggtttca	tcttgcatgt	tgattggatg	gccagaccac	cacgtatagt	tacaaaagtc	9900
atacactgca	cagccacaga	cagcctttcc	cgtaggtcac	agtgcacaat	cttagacctg	9960
ttcacctgca	ggaacctcac	attagaatta	gcagccatga	cccctcatcc	atttatttaag	10020
atcacacctt	ccagagaagc	agtgaacaca	ttaggggccac	acttttcaaa	atagcaattg	10080
actagatttg	accagggtgc	aaatcagatt	ggcaaggatc	tcaaaccctt	cacagaagaa	10140
gaatatctca	tgaaaacaac	aatctctaga	cagaagaaag	ataagatggc	tacataaagt	10200
gatttaggat	gtaaggacca	tctttacata	tttgtatgag	cattaatgca	gaaacatgat	10260
tatagtattt	cattataact	gcagttagca	gaacctagac	aaatggataa	acattacatg	10320
gtgtctccac	catttcta	acgtctctca	aataattaga	acttccataa	gtgaaatggg	10380
tgagttgtga	tgctgagagc	tcccagtcct	tgagcaacct	aactatccct	gagcaagtgt	10440
gttagggacc	tgagcttcat	tataaccagt	cctgacatat	tcatattcat	attcttttga	10500
ggaaaagaaa	aaatcaaaat	gaagtgactt	ccacaacatg	acctgaataa	aaacagatat	10560
ctgtggaaaa	atcaataaat	aataatttcc	atcaagcatg	tctctccagt	gaatcaagag	10620
agataaacctc	attagaacat	tttctctaga	aacataaatt	aaaaaggact	gacagatgga	10680
aataagaaaa	atagcaaatc	tagataggac	tccagggtgaa	tagattttcca	atatttatgc	10740
agagaagttt	tgagagcagg	aaaagtagga	ggaagcagag	gaagacagga	tccaagcttc	10800
ttttcttata	tgattttttc	caaggcctac	attttgttac	tgtttttttg	tgtcaacttg	10860
gctggattgt	gacacctaca	tatttaatac	aacaacaatc	ctgttgtttc	tgtgaaagtg	10920
tcaattgtgt	agttgtgatt	tacatctata	atcagttgac	tttaagaaaa	ggagataatg	10980
ttaaaattata	ataatatggg	tggtctttat	gcaatcacct	gaaggtctta	agagaaaaaa	11040
aactcagggt	tccaaaagaa	aggattctgc	ctcaaggctg	ttatatcaaa	tcttgccctga	11100
ttttccagca	tcccctacag	atttaaaaca	tgccaagacc	cacaactgca	ggagccaatt	11160
cctttaaata	aacctatat	acataatata	cattatatat	gtacatttat	ctatatgtac	11220
atattacata	gccattgcct	aaaataaatt	atattgtgtac	acattttata	tatgcatata	11280
tataaaacat	gcacacacat	acatatatat	ggatgagtc	cctttatact	tggtttccct	11340
ttctgcagtt	ttagtcaccc	acaatcaacc	ccaatccaaa	aatattacag	tatttcaaga	11400
gaaagaagga	tagagagagg	gagattacat	tacatatagat	gtaaatatat	tagagaatat	11460
tgttatagaa	gctctgtttt	attattagtt	ttgttgtaac	tactcacag	tgccataat	11520
aaaaattaaa	gtaatatata	catctatgta	taggaaaaaa	cataaacact	attatcatat	11580
aatttggtac	tatctgtggt	ttcaggcatc	cgtgggggt	catgtgatat	atcccctgtg	11640
cataaggatg	aactaatgta	ttcttttgat	ttgcctattg	ctctgtgtct	ctggagaacc	11700
tggctgacac	agataccagt	caccatgaca	catgccaaag	ttcaagtcac	tgcaaaccta	11760

catctctgtt	tgctctgatt	caaaagagag	catttcacac	acttgctcac	ttggtctgtg	11820
gtcattttcc	tctggagaat	gttttccatc	atccttcaga	cgaatcttca	agtcacttta	11880
gcacaatgct	tgcaacatag	ctcaccctga	ataaagatag	ctcctgtgtt	tataatgact	11940
gcccagaacc	aaaccaggaa	gctgccagaa	gttacaacct	atcagggaca	ctaaaacatc	12000
cctgggataa	aatatggtgc	tggttaactc	agggtgtccac	tcattctctt	accaactagt	12060
gaaaaagaaa	tgcatcccat	gtttaccacg	tagacacagc	ctcagctgga	aatagaagtt	12120
ctcctggagg	gcacccctct	ttctgctgct	gctgagctct	ttttggaaga	ggaacttgca	12180
aactgcatag	tgagctata	cagggaagc	agcaggagga	ccctacccat	ttataggatg	12240
gcctgaatta	gttgagtctg	aaactaaaca	tggtttacca	ggaacagggg	aagaatttta	12300
ttgccgaatg	ttttaagaca	tgtcacaaga	catagtcaat	gtgtgcaaag	tcacatataa	12360
taaatgtgta	ctataaatct	cggctttaca	ccatatagac	aaattttatt	actagaaaat	12420
tatttccact	tcgttaagga	accagaacga	tagttacaga	agcttatttc	aaaggagcac	12480
taatttactt	tatagccttg	ggttttgtcc	cagtactggg	aactaacagt	gctctgactt	12540
cttcagttg	ttggcgggtc	ctggtttgac	catgtgaaag	gatggtgggc	tgcaaaagac	12600
atgcaccgga	tcctctacct	cttctacgag	gatattaaaa	aagtaagtgg	cactgagact	12660
tataggtcag	accagaaac	cctcctgaca	atggttattct	gttaaaaagc	tgtgtcttta	12720
attggccaag	ttcttcttct	ttcctccctc	ttcacaatgc	cttttctctc	catgatcaga	12780
atccaaaaca	tgagatccac	aagggtgttg	aattcttggg	gaaaacttgg	tcagggtgatg	12840
ttataaacia	gattgtccac	catacctcat	ttgatgtaat	gaaggataat	cccatggcca	12900
accatactgc	ggtacctgct	cacatattca	atcactccat	ctcaaaattt	atgaggaaag	12960
gttggtggca	tttcttttcc	ttaactgaac	tctaaaaaat	tttctacctt	atatgctaaa	13020
ataattttca	acctaatttt	caggcagaag	tgactcattt	cagttaaatt	ttgaatctct	13080
gctcccttca	ccttgctgtg	ttgcagacag	ccaatgtcag	tggttctgaa	acttgagtca	13140
cattagaacc	cttcagggcc	ttgctaaagc	tctgattgct	gggtccctct	cagagatact	13200
gattccacag	atccagcagt	agccctcaaa	tttgcttttc	tctcaagtac	tcagggtgatg	13260
ctgatggtgc	tggtcacttt	gattacaata	cccacctcaa	ccatgaactt	ccctttgaag	13320
gcttgtgcat	cctctgagca	gctttgaaca	ctcatcttta	gtctatccct	gtagttcaaa	13380
accctagcta	agcacttagt	acttggattt	gtaactactg	atattcatgt	ctgtctccaa	13440
aataagatga	taggctgtcc	tgaagagagt	gtagtggtca	gttttgttcc	actagaacct	13500
agtatagaga	ctcatacctc	aaaacaactc	agtaatggcc	tggttgtgta	gtgtacagat	13560
gaatgaacat	tatttctgtc	ctcaacaagt	taacattcta	gatacatgca	aaaatagctg	13620
caaaaagtta	taaaacaaga	agtaaagtgg	aagctatact	aggaattccc	taataaccagt	13680
tctcctggct	gtatcaaaat	tacctttaaa	aacagtgacc	ccattccaga	acattccaat	13740
taactagttg	caagctggaa	tctagaattt	gatattatgg	gcaagcattt	cagattaatc	13800
ctcttgtcaa	agggtaggaa	accagtagaa	ataaagtact	aggataaact	agagaaacia	13860
ttcattagta	cagcatttgt	tggtctgaca	aggtagaaca	gtttgcagaa	gatccctagt	13920
atccaaatgt	ctattccagt	ggatttacta	tttaatttta	cccaacaagt	aatatcttct	13980
acaatgaggt	tactgacatc	ttgtaacgtc	tttcaactgtc	cctgggagaa	taagataggc	14040
tgtcccccag	gaagtccatg	atggtaacca	gctgtagact	ttgggttggg	tacactagag	14100
ccaggagtgt	accctggagc	agaagattca	gcacagtggg	gcacctcagt	ggggcctgca	14160
gcatttgagg	aaggagtata	aggatgctgt	gtgccttgtg	gtaagaaaagc	caagtggagt	14220
ggaaatgagg	gaccattcat	agtggaaaaga	ccagagggag	tggtgactgag	ggaccctcac	14280
tcattgggaaa	atcagctggg	taggcctagg	aaccattcac	tatagaaaag	ttaggtggaa	14340
tggtgtcaga	gtctgtcatg	aacttctttg	atgtcctaca	gggatgcctg	gagactggaa	14400
gaaccacttt	actgtggcct	tgaatgagaa	ctttgataag	cattatgaaa	agaagatggc	14460
agggtccaca	ctgaacttct	gcctggagat	ctgagaggaa	caacaacaaa	ctaggtgaca	14520
gagactatgc	caactatttc	gccttttatt	ctggttagca	aggaaactgtg	actgaatgtg	14580
gagcttatga	gcttcagtc	atctcctata	gtgtggctag	tttgcataaa	tattaaaaca	14640
tgattttaaa	tatcaacaaa	ccagttactc	cagtaataaa	aataagagaa	ttagagagca	14700
gagtcgcct	acatgagttt	ttttgtttgt	ttgtttttta	agtacaggta	tgttttattg	14760
tgcatgacag	acagagcaaa	aacaaacaat	tgcatcatgg	attcccatgt	gtgatcccaa	14820
gtagatttca	caagaaaatt	atgcataggt	attacaagcc	ccattgttaa	gagagaatgt	14880
aacagcttga	agtgtacatt	ctatactttt	atgtataaat	aataacttcc	aagagaaaag	14940
agctgataag	tacatttcag	agtaccattt	tctgtaatat	aatgatataa	aaataaatta	15000
ctactgcaaa	atatcaatca	attgcaaaat	gattactgct	ctacttttgg	cttgtaacta	15060
attttctcaa	tcgaaagagt	ttgagttgga	gaaattagtc	agtgagtact	cctgtaaaaa	15120
atcttcccac	gatataaata	aaatgcttaa	tgtatctaat	ctataaattg	agatctggga	15180
caaatgcacc	atgactatgc	attgccattc	tcttaataac	tatgcattgc	catttcttta	15240
ataaagagtc	tctgttgcat	cactctaaca	ataggtatga	cctcagattt	aatataaatt	15300
tagtgcttca	accatgcacc	agtgaagacc	tggtttttta	tatgaccaag	tacaattgtc	15360
ttattagaag	agaaggctct	gaggggctct	agggagtagg	tggtatgcaa	cagggctgat	15420
ggcctcagag	atgacagcat	gtaacatgta	agaaaggggg	gaaatttgga	aagacttaaa	15480
ctcagaaata	aaattaaaaga	agtgtaaaaa	ggtattcagt	taacatcttg	aaaggaatc	15540

aggaacaata	tacatagctc	ccagtacaca	gagcaaaagta	acctccctgt	ggttgtgac	15600
attgtctcta	ccttatcgt	gtgacaaggc	agtgtctattg	ttacgccagg	caaaggggaa	15660
atgacatcct	attctctagt	caactggatg	gaagacaaac	tcaaaggtaa	aaaatagatg	15720
atgagatttt	aagaaaagag	agcagctggt	cctaaggcc	tgctgtaagg	aagcagagag	15780
gatgaagatg	ggggcatttg	aaccagccca	gaggggaccc	tggggtgaaa	gttccccatc	15840
aggcattacc	ccactgccta	catccagcca	gataacccaa	ccacctgaat	taccatcctc	15900
atatttagtt	ggtgatccag	agaagagatt	tctctagtta	ttccctaaag	gtatagcaag	15960
aaaaaaagag	attccttgat	cactcctgcc	tagtcaatgc	actggaagac	tggttcacag	16020
ggctactgtg	ctttgtaatt	ggagacatga	gaatccacaa	tgattagaaa	gctcggggcc	16080
caggtactgt	cagagtcac	agtctacctc	agagaggagg	aggcagatta	aaggaaaagc	16140
aaatttcatt	ttcctactca	gaaatgattt	tctactaatt	gaaaagcaat	tgaatgctgt	16200
caataaagac	atttcctgta	ctaacccttg	actcagagta	ttgatgacca	actatacaaa	16260
gcttatttct	tccatgcaat	ggatgatacc	tgctgcctt	tgaggtatt	aaggggtatg	16320
atgctgagag	tccaggtgta	gaagatggga	tcacgggttt	tgggcaaaagc	attttggcta	16380
tctaagtgtta	agaactgtaa	ggtttgagaa	tgcttgatg	aaagtctcgt	aaagctacaa	16440
acagagttgc	tggtcatttc	tacaaggagg	ctgtgaactg	ctcttcatct	tctagaggca	16500
tattttggct	atgggctact	aagattcaga	caggtgtaag	ataagtttg	ccccatggcc	16560
tccccataga	ctttcccaa	tgtgactggt	cctggactaa	actgagggtc	gggctgctat	16620
ttcctgtggc	ccaataacaa	gatgcagatg	aactggggag	gaagagaagt	tttatttctg	16680
taactgggta	cattcagagg	gcctggaaat	tatcaccaaa	ccaactcaaa	atgacaaaat	16740
ttttcagagc	ttatctacct	tctaagctgt	atgtctacat	gtaagtgtgc	atgccttcta	16800
aagacatatg	tgattaattt	attttaattt	ataactaaga	tctgagtcct	gaagaccttc	16860
ctctgggtgc	taatgaagtt	tgcttaattt	aaatgggtct	ccaggtactg	ggttgcacac	16920
ccttatcttg	tctcctgtta	aactactgag	gtttggggag	ttccttcaga	cctccaataa	16980
acgtgtttgt	ggaggcctgg	ggagtttctt	cagagcccca	ataaaactta	tataatccta	17040
actgggtact	gttaagaact	cctttattat	tttgtcatgt	tgtaaggccc	aggaaacgcc	17100
taggcaaaac	tctggatggg	cttttggtac	atttcagcct	ttgcataagg	gcactggcct	17160
tttttaatat	ttaacttaac	cactcactga	atactgaaac	agttgtgatg	gaggcctgca	17220
ttaatgcaac	ctgcctgcc	caatccccac	tgtaatttg	tgcataattc	ttatcatgct	17280
agtataattga	tttatcatga	gaattgtagg	gatatggggc	attgtaatat	ttctggctac	17340
ttcctgtcta	gtgagtgtca	ttgttatggg	acactgaatg	cagcattggt	ataggggagg	17400
tctatttggt	cccagcagca	ctctttgttt	caggggctta	gaggcagcac	ctgctgaaac	17460
atttagtctt	cagttcacag	ggctttaaga	aagcacaact	taggtttcag	tgatttccag	17520
ttaggaaaaa	tggggtagtg	catgggcttt	catgcagaag	agccttcagt	gcaagtccat	17580
gacaatgttt	gcaactcagt	tttatcctca	aaagctctaa	ctactttttc	ggtattctaa	17640
atctgctttg	ctttaattca	gttttcaaaa	tattcttctt	cctcaggaaa	tagcacatgc	17700
tttatagttg	aatgaatctg	ttggctcctg	gcacttgtaa	gccagatata	tttatctgca	17760
aaatgggtatc	atatctatct	catgagatca	ttgtaaagat	ttacactata	atacttgtaa	17820
agtgtacata	gttgtgttca	agaaatttag	cttcttctc	cctcttgcac	acctcaacc	17880
cacatcataa	agcaaagttt	aatccatata	aacactggta	ggattagtcc	cacatattgg	17940
gacagtggac	tgatttctga	atttcaaacc	atcgtgtctg	agtgtctgca	caaattcact	18000
ctcatgaccc	agtgcacatg	catcctagaa	ttacaaaact	agcatctgag	atgtcctcag	18060
aaatttaatac	aaaagctata	gctttctctc	ttctgtatc	tctttgatca	aaaagatcca	18120
agaaataaag	atctatgctt	ctgggtcaac	ttgagattcc	ttttattcct	cctcttagtg	18180
accttctggt	aagttcatag	gttctttttg	ccatctggtg	tagtccaaat	actagagtta	18240
gttatactat	gcttcaccac	caacattggt	cctccttcac	acagcctttc	ctgttctccc	18300
ccaggagata	tcattgctcc	ttcccaccaa	atctcctggc	actctgtctg	tacgtaggca	18360
gcatgtactt	tctcagccag	ccttgtgatt	gtttgtgttt	atgcttaatt	cctttccttc	18420
aaggtaagcc	tcttgagggc	agggacaacc	ctcatctttg	tggtgctaca	gtatctggga	18480
cagtcacagt	tctggcacta	ggagtcactt	gagaaatatt	tttaaaatgt	gttcaaatga	18540
tcctttaact	catttctctc	caaatactgt	ccaaatggaa	catcatcccc	aagggaagac	18600
ccaacatgat	ttgtttgggt	tgactgttcc	acatacagga	cccaaagcgg	gaaattgaga	18660
agatactgaa	gttcctggaa	aaagacatat	cagaggaaat	tctgaataaa	atcatctatc	18720
acacctcctt	tgatgtaatg	aagcaaaacc	caatgaccaa	ctataccact	ttgccacca	18780
gcattatgga	ccactccatc	tcccctttta	tgaggaaagg	tagataagct	ttgtagtcta	18840
agatgtcaaa	tggaaactctg	tggtccccat	ggtctgctta	gattttccag	taatgtttca	18900
ttctccatta	tttattcttt	ccagcagcac	cactgtacaa	cctttgagag	gcaagttgcc	18960
tgtttctcct	cattcttggt	gggttcctaa	gggtgcatgc	ttacctctcc	ctctactcct	19020
gcagcaatca	ttaagatttt	gccttggttc	agggatgcct	ggagactgga	agaactattt	19080
tactgtggcc	caaaatgaag	aatttgacaa	ggactaccag	aagaagatgg	caggaagcac	19140
cctaaccctt	cgcacagaga	tctgagagca	gtcagggagt	ctgtcctgga	ctttcttacc	19200
agatttttgc	catttgagcc	tcatgatcaa	ggacacttaa	aacaaagaca	cccttccttc	19260
agtctggagc	tgttacacac	tacctgtaga	tgataataac	tcatcaaaat	gtaaccaaat	19320

```

cctgggtaga gtttttaatt aagatacatg atccctcatt tagacaccaa ccacataacg 19380
tactcccctg tcctaaggca aaataaaggc aatttagttc catcactagt ttacaatagt 19440
gaaataaata aagagataaa taagagtaga atttcaatgt aagagaaagt gagcaagaga 19500
agaatgagga tgataagtgg gactaagga taatgtgtgg gggaaactgg ttttattttt 19560
aaaggtcagg aatgagaggc agaggatagt cctactcttg aagatattag attaggaggt 19620
gcaggggggtg gcatgaagga tggaatgctg tgagctgggt ctggctcagg tggcttcaag 19680
ctgaaatggt cccaaaacca aagagaaagg ccacaggaga gcgataagga acacaagatg 19740
aacaagaaac tcagcctact tctgtggcca acacaaggct ggaaagcaga gaaagatttt 19800
atctttcaga tggacttctg agagagagag gaagtcttca gtgagctcaa gggaaaaata 19860
aaataaaaata tttggtaact catggacggt aagaaaagct taaatttcac atctctgtgg 19920
aaaatcggtt ttataacttt ttttaaagg gctcttagca aaactgtcag cattgccaa 19980
ttcttcagct acactaatga ttatgttctt ttcccttctt ttgttaaaac ctgtaccaag 20040
aaaaatatctg ccatcatttt atacaagcta tacaacgac catgtatcat tattctttta 20100
atgtcaggaa ggtacatttt gtgggatagg tggttaactt atcattaac catacaacac 20160
aaaaa                                     20165

```

&lt;210&gt; 8

&lt;211&gt; 283

&lt;212&gt; PRT

&lt;213&gt; HUMAN

&lt;400&gt; 8

```

Glu Val Asn Gly Ile Leu Met Ser Lys Leu Met Ser Asp Asn Trp Asp
 1          5          10          15
Lys Ile Trp Asn Phe Gln Ala Lys Pro Asp Asp Leu Leu Ile Ala Thr
 20          25          30
Tyr Ala Lys Ala Gly Thr Thr Trp Thr Gln Glu Ile Val Asp Met Ile
 35          40          45
Gln Asn Asp Gly Asp Val Gln Lys Cys Gln Arg Ala Asn Thr Tyr Asp
 50          55          60
Arg His Pro Phe Ile Glu Trp Thr Leu Pro Ser Pro Leu Asn Ser Gly
 65          70          75          80
Leu Asp Leu Ala Asn Lys Met Pro Ser Pro Arg Thr Leu Lys Thr His
 85          90          95
Leu Pro Val His Met Leu Pro Pro Ser Phe Trp Lys Glu Asn Ser Lys
100          105          110
Ile Ile Tyr Val Ala Arg Asn Ala Lys Asp Cys Leu Val Ser Tyr Tyr
115          120          125
Tyr Phe Ser Arg Met Asn Lys Met Leu Pro Asp Pro Gly Thr Leu Gly
130          135          140
Glu Tyr Ile Glu Gln Phe Lys Ala Gly Lys Val Leu Trp Gly Ser Trp
145          150          155          160
Tyr Asp His Val Lys Gly Trp Trp Asp Val Lys Asp Gln His Arg Ile
165          170          175
Leu Tyr Leu Phe Tyr Glu Asp Met Lys Glu Asp Pro Lys Arg Glu Ile
180          185          190
Lys Lys Ile Ala Lys Phe Leu Glu Lys Asp Ile Ser Glu Glu Val Leu
195          200          205
Asn Lys Ile Ile Tyr His Thr Ser Phe Asp Val Met Lys Glu Asn Pro
210          215          220
Met Ala Asn Tyr Thr Thr Leu Pro Ser Ser Ile Met Asp His Ser Ile
225          230          235          240
Ser Pro Phe Met Arg Lys Gly Met Pro Gly Asp Trp Lys Asn Tyr Phe
245          250          255
Thr Val Ala Gln Ser Glu Asp Phe Asp Glu Asp Tyr Arg Arg Lys Met
260          265          270
Ala Gly Ser Asn Ile Thr Phe Arg Thr Glu Ile
275          280

```

&lt;210&gt; 9

&lt;211&gt; 283

&lt;212&gt; PRT

&lt;213&gt; HUMAN

&lt;400&gt; 9

Glu Val Asn Gly Ile Leu Met Ser Lys Leu Met Ser Asp Asn Trp Asp  
 1 5 10 15  
 Lys Ile Trp Asn Phe Gln Ala Lys Pro Asp Asp Leu Leu Ile Ala Thr  
 20 25 30  
 Tyr Ala Lys Ala Gly Thr Thr Trp Thr Gln Glu Ile Val Asp Met Ile  
 35 40 45  
 Gln Asn Asp Gly Asp Val Gln Lys Cys Gln Arg Ala Asn Thr Tyr Asp  
 50 55 60  
 Arg His Pro Phe Ile Glu Trp Thr Leu Pro Ser Pro Leu Asn Ser Gly  
 65 70 75 80  
 Leu Asp Leu Ala Asn Lys Met Pro Ser Pro Arg Thr Leu Lys Thr His  
 85 90 95  
 Leu Pro Val His Met Leu Pro Pro Ser Phe Trp Lys Glu Asn Ser Lys  
 100 105 110  
 Ile Ile Tyr Val Ala Arg Asn Ala Lys Asp Cys Leu Val Ser Tyr Tyr  
 115 120 125  
 Tyr Phe Ser Arg Met Asn Lys Met Leu Pro Asp Pro Gly Thr Leu Gly  
 130 135 140  
 Glu Tyr Ile Glu Gln Phe Lys Ala Gly Lys Val Leu Trp Gly Ser Trp  
 145 150 155 160  
 Tyr Asp His Val Lys Gly Trp Trp Asp Val Lys Asp Gln His Arg Ile  
 165 170 175  
 Leu Tyr Leu Phe Tyr Glu Asp Met Lys Glu Asp Pro Lys Arg Glu Ile  
 180 185 190  
 Lys Lys Ile Ala Lys Phe Leu Glu Lys Asp Ile Ser Glu Glu Val Leu  
 195 200 205  
 Asn Lys Ile Ile Tyr His Thr Ser Phe Asp Val Met Lys Glu Asn Pro  
 210 215 220  
 Met Ala Asn Tyr Thr Thr Leu Pro Ser Ser Ile Met Asp His Ser Ile  
 225 230 235 240  
 Ser Pro Phe Met Arg Lys Gly Met Pro Gly Asp Trp Lys Asn Tyr Phe  
 245 250 255  
 Thr Val Ala Gln Ser Glu Asp Phe Asp Glu Asp Tyr Arg Arg Lys Met  
 260 265 270  
 Ala Gly Ser Asn Ile Thr Phe Arg Thr Glu Ile  
 275 280

&lt;210&gt; 10

&lt;211&gt; 283

&lt;212&gt; PRT

&lt;213&gt; HUMAN

&lt;400&gt; 10

Glu Val Asn Gly Ile Leu Met Ser Lys Met Met Ser Glu Asn Trp Asp  
 1 5 10 15  
 Lys Ile Trp Asn Phe Gln Ala Lys Pro Asp Asp Leu Leu Ile Ala Thr  
 20 25 30  
 Tyr Ala Lys Ala Gly Thr Thr Trp Thr Gln Glu Ile Val Asp Met Ile  
 35 40 45  
 Gln Asn Asp Gly Asp Val Gln Lys Cys Gln Arg Ala Asn Thr Tyr Asp  
 50 55 60  
 Arg His Pro Phe Ile Glu Trp Thr Leu Pro Pro Pro Leu Asn Ser Gly  
 65 70 75 80  
 Leu Asp Leu Ala Asn Lys Met Pro Ser Pro Arg Thr Leu Lys Thr His  
 85 90 95  
 Leu Pro Val Gln Met Leu Pro Pro Ser Phe Trp Lys Glu Asn Ser Gln  
 100 105 110  
 Ile Ile Tyr Val Ala Arg Asn Ala Lys Asp Cys Leu Val Ser Tyr Tyr  
 115 120 125

Tyr Phe Ser Arg Met Asn Lys Met Leu Pro Asp Pro Gly Thr Leu Gly  
 130 135 140  
 Glu Tyr Ile Glu Thr Phe Lys Ala Gly Lys Val Leu Trp Gly Ser Trp  
 145 150 155 160  
 Tyr Asp His Val Lys Gly Trp Trp Asp Val Lys Asp Lys His Arg Ile  
 165 170 175  
 Leu Tyr Leu Phe Tyr Glu Asp Met Lys Glu Asp Pro Lys Arg Glu Ile  
 180 185 190  
 Lys Lys Ile Val Lys Phe Leu Glu Lys Asp Ile Ser Glu Glu Val Leu  
 195 200 205  
 Asn Lys Ile Ile His His Thr Ser Phe Asp Val Met Lys Gln Asn Pro  
 210 215 220  
 Met Ala Asn Tyr Thr Thr Leu Pro Ser Ser Ile Met Asp His Ser Ile  
 225 230 235 240  
 Ser Pro Phe Met Arg Lys Gly Met Pro Gly Asp Trp Lys Asn Tyr Phe  
 245 250 255  
 Thr Val Ala Gln Ser Glu Asp Phe Asp Glu Asp Tyr Arg Lys Lys Met  
 260 265 270  
 Ala Gly Ser Thr Ile Thr Phe Arg Thr Glu Ile  
 275 280

<210> 11  
 <211> 52  
 <212> DNA  
 <213> HUMAN

<400> 11  
 ttagtattga tcttaccat cccagattcc cmatggcgaa gattgagaaa aa 52

<210> 12  
 <211> 57  
 <212> DNA  
 <213> HUMAN

<400> 12  
 gcaacattga tctaatttta aagtgtcttc atuttgctgt ctttctcaa aatattc 57

<210> 13  
 <211> 46  
 <212> DNA  
 <213> HUMAN

<400> 13  
 ccaaccctc ctccctgtaa cctctaytct tcctaaaaca caatct 46

<210> 14  
 <211> 49  
 <212> DNA  
 <213> HUMAN

<400> 14  
 agcagccatg acccctcatc catttattma gatcacacct tccagagaa 49

<210> 15  
 <211> 55  
 <212> DNA  
 <213> HUMAN

<400> 15  
 aggaggagc agaggaagac aggawccaag cttcttttct tataatgattt tttcc 55

<210> 16

<211> 52  
<212> DNA  
<213> HUMAN

<400> 16  
ccacttcggtt aaggaaccag aacgawagtt acagaagctt atttcaaagg ag 52

<210> 17  
<211> 62  
<212> DNA  
<213> HUMAN

<400> 17  
ttttaaataat caacaaacca gttactccag yaaataaaat aagagaatta gagagcagag 60  
tc 62

<210> 18  
<211> 42  
<212> DNA  
<213> HUMAN

<400> 18  
aggtattaca agcscattg ttaagagaga atgtaacagc tt 42

<210> 19  
<211> 36  
<212> DNA  
<213> HUMAN

<400> 19  
ttttgtggga taggtgrtta acttatcatt aaacca 36

<210> 20  
<211> 27  
<212> DNA  
<213> HUMAN

<400> 20  
ggtgggaaaa agtmtgtaat ttccaag 27

<210> 21  
<211> 28  
<212> DNA  
<213> HUMAN

<400> 21  
gtgggctgca aaagacaygc accggatc 28

<210> 22  
<211> 25  
<212> DNA  
<213> HUMAN

<400> 22  
ggagaaaact tkgtcaggtg atgtt 25

<210> 23  
<211> 20  
<212> DNA  
<213> HUMAN

<400> 23  
actttactgt ggctwtgaat 20